

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
22 July 2004 (22.07.2004)

PCT

(10) International Publication Number
WO 2004/060278 A2

(51) International Patent Classification⁷: **A61K**
(21) International Application Number:
PCT/US2003/038761

(22) International Filing Date: 5 December 2003 (05.12.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/431,319	6 December 2002 (06.12.2002)	US
10/326,051	18 December 2002 (18.12.2002)	US
10/325,527	18 December 2002 (18.12.2002)	US
10/325,526	18 December 2002 (18.12.2002)	US
10/323,233	18 December 2002 (18.12.2002)	US
60/443,443	29 January 2003 (29.01.2003)	US
60/443,788	30 January 2003 (30.01.2003)	US
60/447,529	14 February 2003 (14.02.2003)	US
60/501,926	11 September 2003 (11.09.2003)	US

(71) Applicant (for all designated States except US): **ISIS PHARMACEUTICALS, INC.** [US/US]; 2292 Faraday Avenue, Carlsbad, CA 92008 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ECKER, David, J.** [US/US]; 1041 Saxony Road, Encinitas, CA 92024 (US). **GRIFFEY, Richard, H.** [US/US]; 360 Barsby Street, Vista, CA 92084 (US). **SAMPATH, Rangarajan** [IN/US]; 12223 Mannix Road, San Diego, CA 92129 (US). **HOFSTADLER, Steven, A.** [US/US]; 5014 Viewridge Way, Oceanside, CA 92056 (US). **MCNEIL,**

John [US/US]; 6389 Castejon Drive, La Jolla, CA 92037 (US). **CROOKE, Stanley, T.** [US/US]; 3211 Piragua Street, Carlsbad, CA 92009 (US). **BLYN, Lawrence, B.** [US/US]; 27892 Perales, Mission Viejo, CA 92692 (US). **RANKEN, Raymond** [US/US]; 231 Horizon Drive, Encinitas, CA 92024 (US). **HALL, Thomas, A.** [US/US]; 5239 Wohlford Street, Oceanside, CA 92056 (US).

(74) Agent: **LEGAARD, Paul, K.**; Cozen O'Connor, 1900 Market Street, Philadelphia, PA 19103 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **METHODS FOR RAPID IDENTIFICATION OF PATHOGENS IN HUMANS AND ANIMALS**

(57) Abstract: The present invention provides methods of: identifying pathogens in biological samples from humans and animals, resolving a plurality of etiologic agents present in samples obtained from humans and animals, determining detailed genetic information about such pathogens or etiologic agents, and rapid detection and identification of bioagents from environmental, clinical or other samples.



WO 2004/060278 A2

METHODS FOR RAPID IDENTIFICATION OF PATHOGENS IN HUMANS AND ANIMALS

FIELD OF THE INVENTION

5 The present invention relates generally to clinical applications of directed to the identification of pathogens in biological samples from humans and animals. The present invention is also directed to the resolution of a plurality of etiologic agents present in samples obtained from humans and animals. The invention is further directed to the determination of detailed genetic information about such pathogens or etiologic agents.

10 The identification of the bioagent is important for determining a proper course of treatment and/or eradication of the bioagent in such cases as biological warfare and natural infections. Furthermore, the determination of the geographic origin of a selected bioagent will facilitate the identification of potential criminal identity. The present invention also relates to methods for rapid detection and identification of bioagents from environmental, clinical or other
15 samples. The methods provide for detection and characterization of a unique base composition signature (BCS) from any bioagent, including bacteria and viruses. The unique BCS is used to rapidly identify the bioagent.

BACKGROUND OF THE INVENTION

20 In the United States, hospitals report well over 5 million cases of recognized infectious disease-related illnesses annually. Significantly greater numbers remain undetected, both in the inpatient and community setting, resulting in substantial morbidity and mortality. Critical intervention for infectious disease relies on rapid, sensitive and specific detection of the offending pathogen, and is central to the mission of microbiology laboratories at medical centers.
25 Unfortunately, despite the recognition that outcomes from infectious illnesses are directly associated with time to pathogen recognition, as well as accurate identification of the class and species of microbe, and ability to identify the presence of drug resistance isolates, conventional hospital laboratories often remain encumbered by traditional slow multi-step culture based assays. Other limitations of the conventional laboratory which have become increasingly
30 apparent include: extremely prolonged wait-times for pathogens with long generation time (up to several weeks); requirements for additional testing and wait times for speciation and identification of antimicrobial resistance; diminished test sensitivity for patients who have received antibiotics; and absolute inability to culture certain pathogens in disease states associated with microbial infection.

For more than a decade, molecular testing has been heralded as the diagnostic tool for the new millennium, whose ultimate potential could include forced obsolescence of traditional hospital laboratories. However, despite the fact that significant advances in clinical application of PCR techniques have occurred, the practicing physician still relies principally on standard techniques. A brief discussion of several existing applications of PCR in the hospital-based setting follows.

Generally speaking molecular diagnostics have been championed for identifying organisms that cannot be grown *in vitro*, or in instances where existing culture techniques are insensitive and/or require prolonged incubation times. PCR-based diagnostics have been successfully developed for a wide variety of microbes. Application to the clinical arena has met with variable success, with only a few assays achieving acceptance and utility.

One of the earliest, and perhaps most widely recognized applications of PCR for clinical practice is in detection of *Mycobacterium tuberculosis*. Clinical characteristics favoring development of a nonculture-based test for tuberculosis include week to month long delays associated with standard testing, occurrence of drug-resistant isolates and public health imperatives associated with recognition, isolation and treatment. Although frequently used as a diagnostic adjunctive, practical and routine clinical application of PCR remains problematic due to significant inter-laboratory variation in sensitivity, and inadequate specificity for use in low prevalence populations, requiring further development at the technical level. Recent advances in the laboratory suggest that identification of drug resistant isolates by amplification of mutations associated with specific antibiotic resistance (e.g., *rpoB* gene in rifampin resistant strains) may be forthcoming for clinical use, although widespread application will require extensive clinical validation.

One diagnostic assay, which has gained widespread acceptance, is for *C. trachomatis*. Conventional detection systems are limiting due to inadequate sensitivity and specificity (direct immunofluorescence or enzyme immunoassay) or the requirement for specialized culture facilities, due to the fastidious characteristics of this microbe. Laboratory development, followed by widespread clinical validation testing in a variety of acute and nonacute care settings have demonstrated excellent sensitivity (90-100%) and specificity (97%) of the PCR assay leading to its commercial development. Proven efficacy of the PCR assay from both genital and urine sampling, have resulted in its application to a variety of clinical setting, most recently including routine screening of patients considered at risk.

While the full potential for PCR diagnostics to provide rapid and critical information to physicians faced with difficult clinical-decisions has yet to be realized, one recently developed

assay provides an example of the promise of this evolving technology. Distinguishing life-threatening causes of fever from more benign causes in children is a fundamental clinical dilemma faced by clinicians, particularly when infections of the central nervous system are being considered. Bacterial causes of meningitis can be highly aggressive, but generally cannot be differentiated on a clinical basis from aseptic meningitis, which is a relatively benign condition that can be managed on an outpatient basis. Existing blood culture methods often take several days to turn positive, and are often confounded by poor sensitivity or false-negative findings in patients receiving empiric antimicrobials. Testing and application of a PCR assay for enteroviral meningitis has been found to be highly sensitive. With reporting of results within 1 day, preliminary clinical trials have shown significant reductions in hospital costs, due to decreased duration of hospital stays and reduction in antibiotic therapy. Other viral PCR assays, now routinely available include those for herpes simplex virus, cytomegalovirus, hepatitis and HIV. Each has a demonstrated cost savings role in clinical practice, including detection of otherwise difficult to diagnose infections and newly realized capacity to monitor progression of disease and response to therapy, vital in the management of chronic infectious diseases.

The concept of a universal detection system has been forwarded for identification of bacterial pathogens, and speaks most directly to the possible clinical implications of a broad-based screening tool for clinical use. Exploiting the existence of highly conserved regions of DNA common to all bacterial species in a PCR assay would empower physicians to rapidly identify the presence of bacteremia, which would profoundly impact patient care. Previous empiric decision making could be abandoned in favor of educated practice, allowing appropriate and expeditious decision-making regarding need for antibiotic therapy and hospitalization.

Experimental work using the conserved features of the 16S rRNA common to almost all bacterial species, is an area of active investigation. Hospital test sites have focused on "high yield" clinical settings where expeditious identification of the presence of systemic bacterial infection has immediate high morbidity and mortality consequences. Notable clinical infections have included evaluation of febrile infants at risk for sepsis, detection of bacteremia in febrile neutropenic cancer patients, and examination of critically ill patients in the intensive care unit. While several of these studies have reported promising results (with sensitivity and specificity well over 90%), significant technical difficulties (described below) remain, and have prevented general acceptance of this assay in clinics and hospitals (which remain dependent on standard blood culture methodologies). Even the revolutionary advances of real-time PCR technique, which offers a quantitative more reproducible and technically simpler system, remains encumbered by inherent technical limitations of the PCR assay.

The principle shortcomings of applying PCR assays to the clinical setting include: inability to eliminate background DNA contamination; interference with the PCR amplification by substrates present in the reaction; and limited capacity to provide rapid reliable speciation, antibiotic resistance and subtype identification. Some laboratories have recently made progress
5 in identifying and removing inhibitors; however background contamination remains problematic, and methods directed towards eliminating exogenous sources of DNA report significant diminution in assay sensitivity. Finally, while product identification and detailed characterization has been achieved using sequencing techniques, these approaches are laborious and time-intensive thus detracting from its clinical applicability.

10 Rapid and definitive microbial identification is desirable for a variety of industrial, medical, environmental, quality, and research reasons. Traditionally, the microbiology laboratory has functioned to identify the etiologic agents of infectious diseases through direct examination and culture of specimens. Since the mid-1980s, researchers have repeatedly demonstrated the practical utility of molecular biology techniques, many of which form the basis of clinical
15 diagnostic assays. Some of these techniques include nucleic acid hybridization analysis, restriction enzyme analysis, genetic sequence analysis, and separation and purification of nucleic acids (See, e.g., J. Sambrook, E. F. Fritsch, and T. Maniatis, *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989). These procedures, in general, are time-consuming and tedious. Another option is the polymerase chain
20 reaction (PCR) or other amplification procedure that amplifies a specific target DNA sequence based on the flanking primers used. Finally, detection and data analysis convert the hybridization event into an analytical result.

Other not yet fully realized applications of PCR for clinical medicine is the identification of infectious causes of disease previously described as idiopathic (e.g. *Bartonella*
25 *henselae* in bacillary angiomatosis, and *Tropheryma whippellii* as the uncultured bacillus associated with Whipple's disease). Further, recent epidemiological studies which suggest a strong association between *Chlamydia pneumonia* and coronary artery disease, serve as example of the possible widespread, yet undiscovered links between pathogen and host which may ultimately allow for new insights into pathogenesis and novel life sustaining or saving
30 therapeutics.

For the practicing clinician, PCR technology offers a yet unrealized potential for diagnostic omnipotence in the arena of infectious disease. A universal reliable infectious disease detection system would certainly become a fundamental tool in the evolving diagnostic armamentarium of the 21st century clinician. For front line emergency physicians, or physicians

working in disaster settings, a quick universal detection system, would allow for molecular triage and early aggressive targeted therapy. Preliminary clinical studies using species specific probes suggest that implementing rapid testing in acute care setting is feasible. Resources could thus be appropriately applied, and patients with suspected infections could rapidly be risk stratified to the different treatment settings, depending on the pathogen and virulence. Furthermore, links with data management systems, locally regionally and nationally, would allow for effective epidemiological surveillance, with obvious benefits for antibiotic selection and control of disease outbreaks.

For the hospitalists, the ability to speciate and subtype would allow for more precise decision-making regarding antimicrobial agents. Patients who are colonized with highly contagious pathogens could be appropriately isolated on entry into the medical setting without delay. Targeted therapy will diminish development of antibiotic resistance. Furthermore, identification of the genetic basis of antibiotic resistant strains would permit precise pharmacologic intervention. Both physician and patient would benefit with less need for repetitive testing and elimination of wait times for test results.

It is certain that the individual patient will benefit directly from this approach. Patients with unrecognized or difficult to diagnose infections would be identified and treated promptly. There will be reduced need for prolonged inpatient stays, with resultant decreases in iatrogenic events.

Mass spectrometry provides detailed information about the molecules being analyzed, including high mass accuracy. It is also a process that can be easily automated. Low-resolution MS may be unreliable when used to detect some known agents, if their spectral lines are sufficiently weak or sufficiently close to those from other living organisms in the sample. DNA chips with specific probes can only determine the presence or absence of specifically anticipated organisms. Because there are hundreds of thousands of species of benign bacteria, some very similar in sequence to threat organisms, even arrays with 10,000 probes lack the breadth needed to detect a particular organism.

Antibodies face more severe diversity limitations than arrays. If antibodies are designed against highly conserved targets to increase diversity, the false alarm problem will dominate, again because threat organisms are very similar to benign ones. Antibodies are only capable of detecting known agents in relatively uncluttered environments.

Several groups have reported detection of PCR products using high resolution electrospray ionization-Fourier transform-ion cyclotron resonance mass spectrometry (ESI-FT-ICR MS). Accurate measurement of exact mass combined with knowledge of the number of at

least one nucleotide allowed calculation of the total base composition for PCR duplex products of approximately 100 base pairs. (Aaserud et al., J. Am. Soc. Mass Spec., 1996, 7, 1266-1269; Muddiman et al., Anal. Chem., 1997, 69, 1543-1549; Wunschel et al., Anal. Chem., 1998, 70, 1203-1207; Muddiman et al., Rev. Anal. Chem., 1998, 17, 1-68). Electrospray ionization-Fourier transform-ion cyclotron resistance (ESI-FT-ICR) MS may be used to determine the mass of double-stranded, 500 base-pair PCR products via the average molecular mass (Hurst et al., Rapid Commun. Mass Spec. 1996, 10, 377-382). Use of matrix-assisted laser desorption ionization-time of flight (MALDI-TOF) mass spectrometry for characterization of PCR products has been described. (Muddiman et al., Rapid Commun. Mass Spec., 1999, 13, 1201-1204). However, the degradation of DNAs over about 75 nucleotides observed with MALDI limited the utility of this method.

U.S. Patent No. 5,849,492 reports a method for retrieval of phylogenetically informative DNA sequences which comprise searching for a highly divergent segment of genomic DNA surrounded by two highly conserved segments, designing the universal primers for PCR amplification of the highly divergent region, amplifying the genomic DNA by PCR technique using universal primers, and then sequencing the gene to determine the identity of the organism.

U.S. Patent No. 5,965,363 reports methods for screening nucleic acids for polymorphisms by analyzing amplified target nucleic acids using mass spectrometric techniques and to procedures for improving mass resolution and mass accuracy of these methods.

WO 99/14375 reports methods, PCR primers and kits for use in analyzing preselected DNA tandem nucleotide repeat alleles by mass spectrometry.

WO 98/12355 reports methods of determining the mass of a target nucleic acid by mass spectrometric analysis, by cleaving the target nucleic acid to reduce its length, making the target single-stranded and using MS to determine the mass of the single-stranded shortened target. Also reported are methods of preparing a double-stranded target nucleic acid for MS analysis comprising amplification of the target nucleic acid, binding one of the strands to a solid support, releasing the second strand and then releasing the first strand which is then analyzed by MS. Kits for target nucleic acid preparation are also provided.

PCT WO97/33000 reports methods for detecting mutations in a target nucleic acid by nonrandomly fragmenting the target into a set of single-stranded nonrandom length fragments and determining their masses by MS.

U.S. Patent No. 5,605,798 reports a fast and highly accurate mass spectrometer-based process for detecting the presence of a particular nucleic acid in a biological sample for diagnostic purposes.

WO 98/21066 reports processes for determining the sequence of a particular target nucleic acid by mass spectrometry. Processes for detecting a target nucleic acid present in a biological sample by PCR amplification and mass spectrometry detection are reported, as are methods for detecting a target nucleic acid in a sample by amplifying the target with primers that
5 contain restriction sites and tags, extending and cleaving the amplified nucleic acid, and detecting the presence of extended product, wherein the presence of a DNA fragment of a mass different from wild-type is indicative of a mutation. Methods of sequencing a nucleic acid via mass spectrometry methods are also reported.

WO 97/37041, WO 99/31278 and U.S. Patent No. 5,547,835 report methods of
10 sequencing nucleic acids using mass spectrometry. U.S. Patent Nos. 5,622,824, 5,872,003 and 5,691,141 report methods, systems and kits for exonuclease-mediated mass spectrometric sequencing.

Thus, there is a need for a method for bioagent detection and identification which is both specific and rapid, and in which no nucleic acid sequencing is required. The present
15 invention addresses this need.

SUMMARY OF THE INVENTION

The present invention is directed towards methods of identifying a pathogen in a biological sample by obtaining nucleic acid from a biological sample, selecting at least one pair
20 of intelligent primers with the capability of amplification of nucleic acid of the pathogen, amplifying the nucleic acid with the primers to obtain at least one amplification product, determining the molecular mass of at least one amplification product from which the pathogen is identified. Further, this invention is directed to methods of epidemic surveillance. By identifying a pathogen from samples acquired from a plurality of geographic locations, the spread of the
25 pathogen to a given geographic location can be determined.

The present invention is also directed to methods of diagnosis of a plurality of etiologic agents of disease in an individual by obtaining a biological sample from an individual, isolating nucleic acid from the biological sample, selecting a plurality of amplification primers with the capability of amplification of nucleic acid of a plurality of etiologic agents of disease, amplifying
30 the nucleic acid with a plurality of primers to obtain a plurality of amplification products corresponding to a plurality of etiologic agents, determining the molecular masses of the plurality of unique amplification products which identify the members of the plurality of etiologic agents.

The present invention is also directed to methods of *in silico* screening of primer sets to be used in identification of a plurality of bioagents by preparing a base composition probability cloud plot from a plurality of base composition signatures of the plurality of bioagents generated *in silico*, inspecting the base composition probability cloud plot for overlap of clouds from
5 different bioagents, and choosing primer sets based on minimal overlap of the clouds.

The present invention is also directed to methods of predicting the identity of a bioagent with a heretofore unknown base composition signature by preparing a base composition probability cloud plot from a plurality of base composition signatures of the plurality of bioagents which includes the heretofore unknown base composition, inspecting the base
10 composition probability cloud for overlap of the heretofore unknown base composition with the cloud of a known bioagent such that overlap predicts that the identity of the bioagent with a heretofore unknown base composition signature equals the identity of the known bioagent.

The present invention is also directed to methods for determining a subspecies characteristic for a given pathogen in a biological sample by identifying the pathogen in a
15 biological sample using broad range survey primers or division-wide primers, selecting at least one pair of drill-down primers to amplify nucleic acid segments which provide a subspecies characteristic about the pathogen, amplifying the nucleic acid segments to produce at least one drill-down amplification product and determining the base composition signature of the drill-down amplification product wherein the base composition signature provides a subspecies
20 characteristic about the pathogen.

The present invention is also directed to methods of pharmacogenetic analysis by obtaining a sample of genomic DNA from an individual, selecting a segment of the genomic DNA which provides pharmacogenetic information, using at least one pair of intelligent primers to produce an amplification product which comprises the segment of genomic DNA and
25 determining the base composition signature of the amplification product, wherein the base composition signature provides pharmacogenetic information about said individual.

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A-1H and Figure 2 are consensus diagrams that show examples of conserved
30 regions from 16S rRNA (Fig. 1A-1, 1A-2, 1A-3, 1A-4, and 1A-5), 23S rRNA (3'-half, Fig. 1B, 1C, and 1D; 5'-half, Fig. 1E-F), 23S rRNA Domain I (Fig. 1G), 23S rRNA Domain IV (Fig. 1H) and 16S rRNA Domain III (Fig. 2) which are suitable for use in the present invention. Lines with arrows are examples of regions to which intelligent primer pairs for PCR are designed. The label for each primer pair represents the starting and ending base number of the amplified region on

the consensus diagram. Bases in capital letters are greater than 95% conserved; bases in lower case letters are 90-95% conserved, filled circles are 80-90% conserved; and open circles are less than 80% conserved. The label for each primer pair represents the starting and ending base number of the amplified region on the consensus diagram. The nucleotide sequence of the 16S rRNA consensus sequence is SEQ ID NO:3 and the nucleotide sequence of the 23S rRNA consensus sequence is SEQ ID NO:4.

Figure 2 shows a typical primer amplified region from the 16S rRNA Domain III shown in Figure 1A-1.

Figure 3 is a schematic diagram showing conserved regions in RNase P. Bases in capital letters are greater than 90% conserved; bases in lower case letters are 80-90% conserved; filled circles designate bases which are 70-80% conserved; and open circles designate bases that are less than 70% conserved.

Figure 4 is a schematic diagram of base composition signature determination using nucleotide analog "tags" to determine base composition signatures.

Figure 5 shows the deconvoluted mass spectra of a *Bacillus anthracis* region with and without the mass tag phosphorothioate A (A*). The two spectra differ in that the measured molecular weight of the mass tag-containing sequence is greater than the unmodified sequence.

Figure 6 shows base composition signature (BCS) spectra from PCR products from *Staphylococcus aureus* (*S. aureus* 16S_1337F) and *Bacillus anthracis* (*B. anthr.* 16S_1337F), amplified using the same primers. The two strands differ by only two (AT-->CG) substitutions and are clearly distinguished on the basis of their BCS.

Figure 7 shows that a single difference between two sequences (A14 in *B. anthracis* vs. A15 in *B. cereus*) can be easily detected using ESI-TOF mass spectrometry.

Figure 8 is an ESI-TOF of *Bacillus anthracis* spore coat protein sspE 56mer plus calibrant. The signals unambiguously identify *B. anthracis* versus other *Bacillus* species.

Figure 9 is an ESI-TOF of a *B. anthracis* synthetic 16S_1228 duplex (reverse and forward strands). The technique easily distinguishes between the forward and reverse strands.

Figure 10 is an ESI-FTICR-MS of a synthetic *B. anthracis* 16S_1337 46 base pair duplex.

Figure 11 is an ESI-TOF-MS of a 56mer oligonucleotide (3 scans) from the *B. anthracis* saspB gene with an internal mass standard. The internal mass standards are designated by asterisks.

Figure 12 is an ESI-TOF-MS of an internal standard with 5 mM TBA-TFA buffer showing that charge stripping with tributylammonium trifluoroacetate reduces the most abundant charge state from $[M-8H+]^8-$ to $[M-3H+]^3-$.

Figure 13 is a portion of a secondary structure defining database according to one embodiment of the present invention, where two examples of selected sequences are displayed graphically thereunder.

Figure 14 is a three dimensional graph demonstrating the grouping of sample molecular weight according to species.

Figure 15 is a three dimensional graph demonstrating the grouping of sample molecular weights according to species of virus and mammal infected.

Figure 16 is a three dimensional graph demonstrating the grouping of sample molecular weights according to species of virus, and animal-origin of infectious agent.

Figure 17 is a figure depicting how a typical triangulation method of the present invention provides for the identification of an unknown bioagent without prior knowledge of the unknown agent. The use of different primer sets to distinguish and identify the unknown is also depicted as primer sets I, II and III within this figure. A three-dimensional graph depicts all of bioagent space (170), including the unknown bioagent, which after use of primer set I (171) according to a method according to the present invention further differentiates and classifies bioagents according to major classifications (176) which, upon further analysis using primer set II (172) differentiates the unknown agent (177) from other, known agents (173) and finally, the use of a third primer set (175) further specifies subgroups within the family of the unknown (174).

Figure 18 shows a representative base composition probability cloud for a region of the RNA polymerase B gene from a cluster of enterobacteria. The dark spheres represent the actual base composition of the organisms. The lighter spheres represent the transitions among base compositions observed in different isolates of the same species of organism.

Figure 19 shows resolution of enterobacteriae members with primers targeting RNA polymerase B (rpoB). A single pair of primers targeting a hyper-variable region within rpoB was sufficient to resolve most members of this group at the genus level (*Salmonella* from *Escherichia* from *Yersinia*) as well as the species/strain level (*E. coli* K12 from O157). All organisms with the exception of *Y. pestis* were tested in the lab and the measured base counts (shown with arrow) matched the predictions in every case.

Figure 20 shows detection of *S. aureus* in blood. Spectra on the right indicate signals corresponding to *S. aureus* detection in spiked wells A1 and A4 with no detection in control wells A2 and A3.

Figure 21 shows a representative base composition distribution of human adenovirus strain types for a single primer pair region on the hexon gene. The circles represent different adenovirus sequences in our database that were used for primer design. Measurement of masses and base counts for each of the unknown samples A, B, C and D matched one or more of the known groups of adenoviruses.

Figure 22 shows a representative broad range survey/drill-down process as applied to emm-typing of *streptococcus pyogenes* (Group A *Streptococcus*: GAS). Genetic material is extracted (201) and amplified using broad range survey primers (202). The amplification products are analyzed (203) to determine the presence and identity of bioagents at the species level. If *Streptococcus pyogenes* is detected (204), the emm-typing "drill-down" primers are used to reexamine the extract to identify the emm-type of the sample (205). Different sets of drill down primers can be employed to determine a subspecies characteristic for various strains of various bioagents (206).

Figure 23 shows a representative base composition distribution of bioagents detected in throat swabs from military personnel using a broad range primer pair directed to 16S rRNA.

Figure 24 shows a representative deconvoluted ESI-FTICR spectra of the PCR products produced by the gtr primer for samples 12 (top) and 10 (bottom) corresponding to emm types 3 and 6, respectively. Accurate mass measurements were obtained by using an internal mass standard and post-calibrating each spectrum; the experimental mass measurement uncertainty on each strand is + 0.035 Daltons (1 ppm). Unambiguous base compositions of the amplicons were determined by calculating all putative base compositions of each stand within the measured mass (and measured mass uncertainty) and selecting complementary pairs within the mass measurement uncertainty. In all cases there was only one base composition within 25 ppm. The measured mass difference of 15.985 Da between the strands shown on the left is in excellent agreement with the theoretical mass difference of 15.994 Da expected for an A to G substitution.

Figure 25 shows representative results of the base composition analysis on throat swab samples using the six primer pairs, 5'-emm gene sequencing and the MLST gene sequencing method of the present invention for an outbreak of *Streptococcus pyogenes* (group A *streptococcus*; GAS) at a military training camp.

Figure 26 shows: a) a representative ESI-FTICR mass spectrum of a restriction digest of a 986 bp region of the 16S ribosomal gene from *E. coli* K12 digested with a mixture of *Bst*NI,

BsmFI, *BfaI*, and *NcoI*; b) a deconvoluted representation (neutral mass) of the above spectrum showing the base compositions derived from accurate mass measurements of each fragment; and c) a representative reconstructed restriction map showing complete base composition coverage for nucleotides 1-856. The *NcoI* did not cut.

- 5 Figure 27 shows a representative base composition distribution of *poxviruses* for a single primer pair region on the DNA-dependent polymerase B gene (DdDpB). The spheres represent different *poxvirus* sequences that were used for primer design.

DESCRIPTION OF EMBODIMENTS

- 10 The present invention provides, *inter alia*, methods for detection and identification of bioagents in an unbiased manner using "bioagent identifying amplicons." "Intelligent primers" are selected to hybridize to conserved sequence regions of nucleic acids derived from a bioagent and which bracket variable sequence regions to yield a bioagent identifying amplicon which can be amplified and which is amenable to molecular mass determination. The molecular mass then
15 provides a means to uniquely identify the bioagent without a requirement for prior knowledge of the possible identity of the bioagent. The molecular mass or corresponding "base composition signature" (BCS) of the amplification product is then matched against a database of molecular masses or base composition signatures. Furthermore, the method can be applied to rapid parallel "multiplex" analyses, the results of which can be employed in a triangulation identification
20 strategy. The present method provides rapid throughput and does not require nucleic acid sequencing of the amplified target sequence for bioagent detection and identification.

- In the context of this invention, a "bioagent" is any organism, cell, or virus, living or dead, or a nucleic acid derived from such an organism, cell or virus. Examples of bioagents include, but are not limited, to cells (including, but not limited to, human clinical samples,
25 bacterial cells and other pathogens) viruses, fungi, and protists, parasites, and pathogenicity markers (including, but not limited to, pathogenicity islands, antibiotic resistance genes, virulence factors, toxin genes and other bioregulating compounds). Samples may be alive or dead or in a vegetative state (for example, vegetative bacteria or spores) and may be encapsulated or bioengineered. In the context of this invention, a "pathogen" is a bioagent that
30 causes a disease or disorder.

Despite enormous biological diversity, all forms of life on earth share sets of essential, common features in their genomes. Bacteria, for example have highly conserved sequences in a variety of locations on their genomes. Most notable is the universally conserved region of the ribosome, but there are also conserved elements in other non-coding RNAs, including RNase P

and the signal recognition particle (SRP) among others. Bacteria have a common set of absolutely required genes. About 250 genes are present in all bacterial species (Mushegian et al., Proc. Natl. Acad. Sci. U.S.A., 1996, 93, 10268; and Fraser et al., Science, 1995, 270, 397), including tiny genomes like *Mycoplasma*, *Ureaplasma* and *Rickettsia*. These genes encode proteins involved in translation, replication, recombination and repair, transcription, nucleotide metabolism, amino acid metabolism, lipid metabolism, energy generation, uptake, secretion and the like. Examples of these proteins are DNA polymerase III beta, elongation factor TU, heat shock protein groEL, RNA polymerase beta, phosphoglycerate kinase, NADH dehydrogenase, DNA ligase, DNA topoisomerase and elongation factor G. Operons can also be targeted using the present method. One example of an operon is the bfp operon from enteropathogenic *E. coli*. Multiple core chromosomal genes can be used to classify bacteria at a genus or genus species level to determine if an organism has threat potential. The methods can also be used to detect pathogenicity markers (plasmid or chromosomal) and antibiotic resistance genes to confirm the threat potential of an organism and to direct countermeasures.

Since genetic data provide the underlying basis for identification of bioagents by the methods of the present invention, it is prudent to select segments of nucleic acids which ideally provide enough variability to distinguish each individual bioagent and whose molecular mass is amenable to molecular mass determination. In one embodiment of the present invention, at least one polynucleotide segment is amplified to facilitate detection and analysis in the process of identifying the bioagent. Thus, the nucleic acid segments that provide enough variability to distinguish each individual bioagent and whose molecular masses are amenable to molecular mass determination are herein described as "bioagent identifying amplicons." The term "amplicon" as used herein, refers to a segment of a polynucleotide which is amplified in an amplification reaction. In some embodiments of the present invention, bioagent identifying amplicons comprise from about 45 to about 150 nucleobases (i.e. from about 45 to about 150 linked nucleosides). One of ordinary skill in the art will appreciate that the invention embodies compounds of 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, and 150 nucleobases in length.

As used herein, "intelligent primers" are primers that are designed to bind to highly conserved sequence regions that flank an intervening variable region and yield amplification

products which ideally provide enough variability to distinguish each individual bioagent, and which are amenable to molecular mass analysis. By the term "highly conserved," it is meant that the sequence regions exhibit between about 80-100%, or between about 90-100%, or between about 95-100% identity. The molecular mass of a given amplification product provides a means
5 of identifying the bioagent from which it was obtained, due to the variability of the variable region. Thus, design of intelligent primers involves selection of a variable region with appropriate variability to resolve the identity of a particular bioagent. It is the combination of the portion of the bioagent nucleic acid molecule sequence to which the intelligent primers hybridize and the intervening variable region that makes up the bioagent identifying amplicon. Alternately,
10 it is the intervening variable region by itself that makes up the bioagent identifying amplicon.

It is understood in the art that the sequence of a primer need not be 100% complementary to that of its target nucleic acid to be specifically hybridizable. Moreover, a primer may hybridize over one or more segments such that intervening or adjacent segments are not involved in the hybridization event (e.g., a loop structure or hairpin structure). The primers of
15 the present invention can comprise at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% sequence complementarity to the target region within the highly conserved region to which they are targeted. For example, an intelligent primer wherein 18 of 20 nucleobases are complementary to a highly conserved region would represent 90 percent complementarity to the highly conserved region. In this example, the remaining
20 noncomplementary nucleobases may be clustered or interspersed with complementary nucleobases and need not be contiguous to each other or to complementary nucleobases. As such, a primer which is 18 nucleobases in length having 4 (four) noncomplementary nucleobases which are flanked by two regions of complete complementarity with the highly conserved region would have 77.8% overall complementarity with the highly conserved region and would thus fall
25 within the scope of the present invention. Percent complementarity of a primer with a region of a target nucleic acid can be determined routinely using BLAST programs (basic local alignment search tools) and PowerBLAST programs known in the art (Altschul et al., J. Mol. Biol., 1990, 215, 403-410; Zhang and Madden, Genome Res., 1997, 7, 649-656).

Percent homology, sequence identity or complementarity, can be determined by, for
30 example, the Gap program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, Madison WI), using default settings, which uses the algorithm of Smith and Waterman (Adv. Appl. Math., 1981, 2, 482-489). In some embodiments, complementarity of intelligent primers, is between about 70% and about 80%. In other embodiments, homology, sequence identity or complementarity, is between about 80% and

about 90%. In yet other embodiments, homology, sequence identity or complementarity, is about 90%, about 92%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100%.

The intelligent primers of this invention comprise from about 12 to about 35
5 nucleobases (i.e. from about 12 to about 35 linked nucleosides). One of ordinary skill in the art will appreciate that the invention embodies compounds of 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, or 35 nucleobases in length.

One having skill in the art armed with the preferred bioagent identifying amplicons defined by the primers illustrated herein will be able, without undue experimentation, to identify
10 additional intelligent primers.

In one embodiment, the bioagent identifying amplicon is a portion of a ribosomal RNA (rRNA) gene sequence. With the complete sequences of many of the smallest microbial genomes now available, it is possible to identify a set of genes that defines "minimal life" and identify composition signatures that uniquely identify each gene and organism. Genes that encode core
15 life functions such as DNA replication, transcription, ribosome structure, translation, and transport are distributed broadly in the bacterial genome and are suitable regions for selection of bioagent identifying amplicons. Ribosomal RNA (rRNA) genes comprise regions that provide useful base composition signatures. Like many genes involved in core life functions, rRNA genes contain sequences that are extraordinarily conserved across bacterial domains interspersed
20 with regions of high variability that are more specific to each species. The variable regions can be utilized to build a database of base composition signatures. The strategy involves creating a structure-based alignment of sequences of the small (16S) and the large (23S) subunits of the rRNA genes. For example, there are currently over 13,000 sequences in the ribosomal RNA database that has been created and maintained by Robin Gutell, University of Texas at Austin,
25 and is publicly available on the Institute for Cellular and Molecular Biology web page on the world wide web of the Internet at, for example, "rna.icmb.utexas.edu/." There is also a publicly available rRNA database created and maintained by the University of Antwerp, Belgium on the world wide web of the Internet at, for example, "rna.uia.ac.be."

These databases have been analyzed to determine regions that are useful as bioagent
30 identifying amplicons. The characteristics of such regions include: a) between about 80 and 100%, or greater than about 95% identity among species of the particular bioagent of interest, of upstream and downstream nucleotide sequences which serve as sequence amplification primer sites; b) an intervening variable region which exhibits no greater than about 5% identity among

species; and c) a separation of between about 30 and 1000 nucleotides, or no more than about 50-250 nucleotides, or no more than about 60-100 nucleotides, between the conserved regions.

As a non-limiting example, for identification of *Bacillus* species, the conserved sequence regions of the chosen bioagent identifying amplicon must be highly conserved among
5 all *Bacillus* species while the variable region of the bioagent identifying amplicon is sufficiently variable such that the molecular masses of the amplification products of all species of *Bacillus* are distinguishable.

Bioagent identifying amplicons amenable to molecular mass determination are either of a length, size or mass compatible with the particular mode of molecular mass determination or
10 compatible with a means of providing a predictable fragmentation pattern in order to obtain predictable fragments of a length compatible with the particular mode of molecular mass determination. Such means of providing a predictable fragmentation pattern of an amplification product include, but are not limited to, cleavage with restriction enzymes or cleavage primers, for example.

15 Identification of bioagents can be accomplished at different levels using intelligent primers suited to resolution of each individual level of identification. "Broad range survey" intelligent primers are designed with the objective of identifying a bioagent as a member of a particular division of bioagents. A "bioagent division" is defined as group of bioagents above the species level and includes but is not limited to: orders, families, classes, clades, genera or other
20 such groupings of bioagents above the species level. As a non-limiting example, members of the *Bacillus/Clostridia* group or gamma-proteobacteria group may be identified as such by employing broad range survey intelligent primers such as primers that target 16S or 23S ribosomal RNA.

In some embodiments, broad range survey intelligent primers are capable of
25 identification of bioagents at the species level. One main advantage of the detection methods of the present invention is that the broad range survey intelligent primers need not be specific for a particular bacterial species, or even genus, such as *Bacillus* or *Streptomyces*. Instead, the primers recognize highly conserved regions across hundreds of bacterial species including, but not limited to, the species described herein. Thus, the same broad range survey intelligent primer
30 pair can be used to identify any desired bacterium because it will bind to the conserved regions that flank a variable region specific to a single species, or common to several bacterial species, allowing unbiased nucleic acid amplification of the intervening sequence and determination of its molecular weight and base composition. For example, the 16S_971-1062, 16S_1228-1310 and 16S_1100-1188 regions are 98-99% conserved in about 900 species of bacteria (16S=16S rRNA,

numbers indicate nucleotide position). In one embodiment of the present invention, primers used in the present method bind to one or more of these regions or portions thereof.

Due to their overall conservation, the flanking rRNA primer sequences serve as good intelligent primer binding sites to amplify the nucleic acid region of interest for most, if not all, bacterial species. The intervening region between the sets of primers varies in length and/or composition, and thus provides a unique base composition signature. Examples of intelligent primers that amplify regions of the 16S and 23S rRNA are shown in Figures 1A-1H. A typical primer amplified region in 16S rRNA is shown in Figure 2. The arrows represent primers that bind to highly conserved regions that flank a variable region in 16S rRNA domain III. The amplified region is the stem-loop structure under "1100-1188." It is advantageous to design the broad range survey intelligent primers to minimize the number of primers required for the analysis, and to allow detection of multiple members of a bioagent division using a single pair of primers. The advantage of using broad range survey intelligent primers is that once a bioagent is broadly identified, the process of further identification at species and sub-species levels is facilitated by directing the choice of additional intelligent primers.

"Division-wide" intelligent primers are designed with an objective of identifying a bioagent at the species level. As a non-limiting example, a *Bacillus anthracis*, *Bacillus cereus* and *Bacillus thuringiensis* can be distinguished from each other using division-wide intelligent primers. Division-wide intelligent primers are not always required for identification at the species level because broad range survey intelligent primers may provide sufficient identification resolution to accomplishing this identification objective.

"Drill-down" intelligent primers are designed with an objective of identifying a sub-species characteristic of a bioagent. A "sub-species characteristic" is defined as a property imparted to a bioagent at the sub-species level of identification as a result of the presence or absence of a particular segment of nucleic acid. Such sub-species characteristics include, but are not limited to, strains, sub-types, pathogenicity markers such as antibiotic resistance genes, pathogenicity islands, toxin genes and virulence factors. Identification of such sub-species characteristics is often critical for determining proper clinical treatment of pathogen infections.

Chemical Modifications of Intelligent Primers

Ideally, intelligent primer hybridization sites are highly conserved in order to facilitate the hybridization of the primer. In cases where primer hybridization is less efficient due to lower levels of conservation of sequence, intelligent primers can be chemically modified to improve the efficiency of hybridization.

For example, because any variation (due to codon wobble in the 3rd position) in these conserved regions among species is likely to occur in the third position of a DNA triplet, oligonucleotide primers can be designed such that the nucleotide corresponding to this position is a base which can bind to more than one nucleotide, referred to herein as a "universal base." For example, under this "wobble" pairing, inosine (I) binds to U, C or A; guanine (G) binds to U or C, and uridine (U) binds to U or C. Other examples of universal bases include nitroindoles such as 5-nitroindole or 3-nitropyrrole (Loakes et al., *Nucleosides and Nucleotides*, 1995, 14, 1001-1003), the degenerate nucleotides dP or dK (Hill *et al.*), an acyclic nucleoside analog containing 5-nitroindazole (Van Aerschot et al., *Nucleosides and Nucleotides*, 1995, 14, 1053-1056) or the purine analog 1-(2-deoxy- β -D-ribofuranosyl)-imidazole-4-carboxamide (Sala et al., *Nucl. Acids Res.*, 1996, 24, 3302-3306).

In another embodiment of the invention, to compensate for the somewhat weaker binding by the "wobble" base, the oligonucleotide primers are designed such that the first and second positions of each triplet are occupied by nucleotide analogs which bind with greater affinity than the unmodified nucleotide. Examples of these analogs include, but are not limited to, 2,6-diaminopurine which binds to thymine, propyne T which binds to adenine and propyne C and phenoxazines, including G-clamp, which binds to G. Propynylated pyrimidines are described in U.S. Patent Nos. 5,645,985, 5,830,653 and 5,484,908, each of which is commonly owned and incorporated herein by reference in its entirety. Propynylated primers are claimed in U.S. Serial No. 10/294,203 which is also commonly owned and incorporated herein by reference in entirety. Phenoxazines are described in U.S. Patent Nos. 5,502,177, 5,763,588, and 6,005,096, each of which is incorporated herein by reference in its entirety. G-clamps are described in U.S. Patent Nos. 6,007,992 and 6,028,183, each of which is incorporated herein by reference in its entirety.

A theoretically ideal bioagent detector would identify, quantify, and report the complete nucleic acid sequence of every bioagent that reached the sensor. The complete sequence of the nucleic acid component of a pathogen would provide all relevant information about the threat, including its identity and the presence of drug-resistance or pathogenicity markers. This ideal has not yet been achieved. However, the present invention provides a straightforward strategy for obtaining information with the same practical value based on analysis of bioagent identifying amplicons by molecular mass determination.

In some cases, a molecular mass of a given bioagent identifying amplicon alone does not provide enough resolution to unambiguously identify a given bioagent. For example, the molecular mass of the bioagent identifying amplicon obtained using the intelligent primer pair "16S_971" would be 55622 Da for both *E. coli* and *Salmonella typhimurium*. However, if

additional intelligent primers are employed to analyze additional bioagent identifying amplicons, a "triangulation identification" process is enabled. For example, the "16S_1100" intelligent primer pair yields molecular masses of 55009 and 55005 Da for *E. coli* and *Salmonella typhimurium*, respectively. Furthermore, the "23S_855" intelligent primer pair yields molecular masses of 42656 and 42698 Da for *E. coli* and *Salmonella typhimurium*, respectively. In this basic example, the second and third intelligent primer pairs provided the additional "fingerprinting" capability or resolution to distinguish between the two bioagents.

In another embodiment, the triangulation identification process is pursued by measuring signals from a plurality of bioagent identifying amplicons selected within multiple core genes.

10 This process is used to reduce false negative and false positive signals, and enable reconstruction of the origin of hybrid or otherwise engineered bioagents. In this process, after identification of multiple core genes, alignments are created from nucleic acid sequence databases. The alignments are then analyzed for regions of conservation and variation, and bioagent identifying amplicons are selected to distinguish bioagents based on specific genomic differences. For

15 example, identification of the three part toxin genes typical of *B. anthracis* (Bowen et al., J. Appl. Microbiol., 1999, 87, 270-278) in the absence of the expected signatures from the *B. anthracis* genome would suggest a genetic engineering event.

The triangulation identification process can be pursued by characterization of bioagent identifying amplicons in a massively parallel fashion using the polymerase chain reaction (PCR),

20 such as multiplex PCR, and mass spectrometric (MS) methods. Sufficient quantities of nucleic acids should be present for detection of bioagents by MS. A wide variety of techniques for preparing large amounts of purified nucleic acids or fragments thereof are well known to those of skill in the art. PCR requires one or more pairs of oligonucleotide primers that bind to regions which flank the target sequence(s) to be amplified. These primers prime synthesis of a different

25 strand of DNA with synthesis occurring in the direction of one primer towards the other primer. The primers, DNA to be amplified, a thermostable DNA polymerase (e.g. *Taq* polymerase), the four deoxynucleotide triphosphates, and a buffer are combined to initiate DNA synthesis. The solution is denatured by heating, then cooled to allow annealing of newly added primer, followed by another round of DNA synthesis. This process is typically repeated for about 30 cycles,

30 resulting in amplification of the target sequence.

Although the use of PCR is suitable, other nucleic acid amplification techniques may also be used, including ligase chain reaction (LCR) and strand displacement amplification (SDA). The high-resolution MS technique allows separation of bioagent spectral lines from background spectral lines in highly cluttered environments.

In another embodiment, the detection scheme for the PCR products generated from the bioagent(s) incorporates at least three features. First, the technique simultaneously detects and differentiates multiple (generally about 6-10) PCR products. Second, the technique provides a molecular mass that uniquely identifies the bioagent from the possible primer sites. Finally, the
5 detection technique is rapid, allowing multiple PCR reactions to be run in parallel.

Mass spectrometry (MS)-based detection of PCR products provides a means for determination of BCS that has several advantages. MS is intrinsically a parallel detection scheme without the need for radioactive or fluorescent labels, since every amplification product is identified by its molecular mass. The current state of the art in mass spectrometry is such that
10 less than femtomole quantities of material can be readily analyzed to afford information about the molecular contents of the sample. An accurate assessment of the molecular mass of the material can be quickly obtained, irrespective of whether the molecular weight of the sample is several hundred, or in excess of one hundred thousand atomic mass units (amu) or Daltons. Intact molecular ions can be generated from amplification products using one of a variety of
15 ionization techniques to convert the sample to gas phase. These ionization methods include, but are not limited to, electrospray ionization (ESI), matrix-assisted laser desorption ionization (MALDI) and fast atom bombardment (FAB). For example, MALDI of nucleic acids, along with examples of matrices for use in MALDI of nucleic acids, are described in WO 98/54751 (Genetrace, Inc.).

20 In some embodiments, large DNAs and RNAs, or large amplification products therefrom, can be digested with restriction endonucleases prior to ionization. Thus, for example, an amplification product that was 10 kDa could be digested with a series of restriction endonucleases to produce a panel of, for example, 100 Da fragments. Restriction endonucleases and their sites of action are well known to the skilled artisan. In this manner, mass spectrometry
25 can be performed for the purposes of restriction mapping.

Upon ionization, several peaks are observed from one sample due to the formation of ions with different charges. Averaging the multiple readings of molecular mass obtained from a single mass spectrum affords an estimate of molecular mass of the bioagent. Electrospray ionization mass spectrometry (ESI-MS) is particularly useful for very high molecular weight
30 polymers such as proteins and nucleic acids having molecular weights greater than 10 kDa, since it yields a distribution of multiply-charged molecules of the sample without causing a significant amount of fragmentation.

The mass detectors used in the methods of the present invention include, but are not limited to, Fourier transform ion cyclotron resonance mass spectrometry (FT-ICR-MS), ion trap, quadrupole, magnetic sector, time of flight (TOF), Q-TOF, and triple quadrupole.

In general, the mass spectrometric techniques which can be used in the present invention include, but are not limited to, tandem mass spectrometry, infrared multiphoton dissociation and pyrolytic gas chromatography mass spectrometry (PGC-MS). In one embodiment of the invention, the bioagent detection system operates continually in bioagent detection mode using pyrolytic GC-MS without PCR for rapid detection of increases in biomass (for example, increases in fecal contamination of drinking water or of germ warfare agents). To achieve minimal latency, a continuous sample stream flows directly into the PGC-MS combustion chamber. When an increase in biomass is detected, a PCR process is automatically initiated. Bioagent presence produces elevated levels of large molecular fragments from, for example, about 100-7,000 Da which are observed in the PGC-MS spectrum. The observed mass spectrum is compared to a threshold level and when levels of biomass are determined to exceed a predetermined threshold, the bioagent classification process described hereinabove (combining PCR and MS, such as FT-ICR MS) is initiated. Optionally, alarms or other processes (halting ventilation flow, physical isolation) are also initiated by this detected biomass level.

The accurate measurement of molecular mass for large DNAs is limited by the adduction of cations from the PCR reaction to each strand, resolution of the isotopic peaks from natural abundance ^{13}C and ^{15}N isotopes, and assignment of the charge state for any ion. The cations are removed by in-line dialysis using a flow-through chip that brings the solution containing the PCR products into contact with a solution containing ammonium acetate in the presence of an electric field gradient orthogonal to the flow. The latter two problems are addressed by operating with a resolving power of $>100,000$ and by incorporating isotopically depleted nucleotide triphosphates into the DNA. The resolving power of the instrument is also a consideration. At a resolving power of 10,000, the modeled signal from the $[\text{M}-14\text{H}]^{14+}$ charge state of an 84mer PCR product is poorly characterized and assignment of the charge state or exact mass is impossible. At a resolving power of 33,000, the peaks from the individual isotopic components are visible. At a resolving power of 100,000, the isotopic peaks are resolved to the baseline and assignment of the charge state for the ion is straightforward. The ^{13}C , ^{15}N -depleted triphosphates are obtained, for example, by growing microorganisms on depleted media and harvesting the nucleotides (Batey et al., Nucl. Acids Res., 1992, 20, 4515-4523).

While mass measurements of intact nucleic acid regions are believed to be adequate to determine most bioagents, tandem mass spectrometry (MS^n) techniques may provide more

definitive information pertaining to molecular identity or sequence. Tandem MS involves the coupled use of two or more stages of mass analysis where both the separation and detection steps are based on mass spectrometry. The first stage is used to select an ion or component of a sample from which further structural information is to be obtained. The selected ion is then fragmented
5 using, e.g., blackbody irradiation, infrared multiphoton dissociation, or collisional activation. For example, ions generated by electrospray ionization (ESI) can be fragmented using IR multiphoton dissociation. This activation leads to dissociation of glycosidic bonds and the phosphate backbone, producing two series of fragment ions, called the *w*-series (having an intact 3' terminus and a 5' phosphate following internal cleavage) and the *a*-Base series (having an
10 intact 5' terminus and a 3' furan).

The second stage of mass analysis is then used to detect and measure the mass of these resulting fragments of product ions. Such ion selection followed by fragmentation routines can be performed multiple times so as to essentially completely dissect the molecular sequence of a sample.

15 If there are two or more targets of similar molecular mass, or if a single amplification reaction results in a product that has the same mass as two or more bioagent reference standards, they can be distinguished by using mass-modifying "tags." In this embodiment of the invention, a nucleotide analog or "tag" is incorporated during amplification (e.g., a 5-(trifluoromethyl) deoxythymidine triphosphate) which has a different molecular weight than the unmodified base
20 so as to improve distinction of masses. Such tags are described in, for example, PCT WO97/33000, which is incorporated herein by reference in its entirety. This further limits the number of possible base compositions consistent with any mass. For example, 5-(trifluoromethyl)deoxythymidine triphosphate can be used in place of dTTP in a separate nucleic acid amplification reaction. Measurement of the mass shift between a conventional amplification
25 product and the tagged product is used to quantitate the number of thymidine nucleotides in each of the single strands. Because the strands are complementary, the number of adenosine nucleotides in each strand is also determined.

In another amplification reaction, the number of G and C residues in each strand is determined using, for example, the cytidine analog 5-methylcytosine (5-meC) or propyne C. The
30 combination of the A/T reaction and G/C reaction, followed by molecular weight determination, provides a unique base composition. This method is summarized in Figure 4 and Table 1.

Table 1

Mass tag	Double strand sequence	Single strand Sequence	Total mass this strand	Base info this strand	Base info other strand	Total base comp. Top strand	Total base comp. Bottom strand
T* _{mass} (T*-T) = x	T*ACGT*ACGT* AT*GCAT*GCA	T*ACGT*ACGT*	3x	3T	3A	3T 2A 2C 2G	3A 2T 2G 2C
		AT*GCAT*GCA	2x	2T	2A		
C* _{mass} (C*-C) = y	TAC*GTAC*GT ATGC*ATGC*A	TAC*GTAC*GT	2x	2C	2G		
		ATGC*ATGC*A	2x	2C	2G		

The mass tag phosphorothioate A (A*) was used to distinguish a *Bacillus anthracis* cluster. The *B. anthracis* (A₁₄G₉C₁₄T₉) had an average MW of 14072.26, and the *B. anthracis* 5 (A₁A*₁₃G₉C₁₄T₉) had an average molecular weight of 14281.11 and the phosphorothioate A had an average molecular weight of +16.06 as determined by ESI-TOF MS. The deconvoluted spectra are shown in Figure 5.

In another example, assume the measured molecular masses of each strand are 30,000.115Da and 31,000.115 Da respectively, and the measured number of dT and dA residues 10 are (30,28) and (28,30). If the molecular mass is accurate to 100 ppm, there are 7 possible combinations of dG+dC possible for each strand. However, if the measured molecular mass is accurate to 10 ppm, there are only 2 combinations of dG+dC, and at 1 ppm accuracy there is only one possible base composition for each strand.

Signals from the mass spectrometer may be input to a maximum-likelihood detection 15 and classification algorithm such as is widely used in radar signal processing. The detection processing uses matched filtering of BCS observed in mass-basecount space and allows for detection and subtraction of signatures from known, harmless organisms, and for detection of unknown bioagent threats. Comparison of newly observed bioagents to known bioagents is also possible, for estimation of threat level, by comparing their BCS to those of known organisms and 20 to known forms of pathogenicity enhancement, such as insertion of antibiotic resistance genes or toxin genes.

Processing may end with a Bayesian classifier using log likelihood ratios developed from the observed signals and average background levels. The program emphasizes performance predictions culminating in probability-of-detection versus probability-of-false-alarm plots for conditions involving complex backgrounds of naturally occurring organisms and environmental
5 contaminants. Matched filters consist of a priori expectations of signal values given the set of primers used for each of the bioagents. A genomic sequence database (e.g. GenBank) is used to define the mass basecount matched filters. The database contains known threat agents and benign background organisms. The latter is used to estimate and subtract the signature produced by the background organisms. A maximum likelihood detection of known background organisms is
10 implemented using matched filters and a running-sum estimate of the noise covariance.

Background signal strengths are estimated and used along with the matched filters to form signatures that are then subtracted. The maximum likelihood process is applied to this "cleaned up" data in a similar manner employing matched filters for the organisms and a running-sum estimate of the noise-covariance for the cleaned up data.

15 Although the molecular mass of amplification products obtained using intelligent primers provides a means for identification of bioagents, conversion of molecular mass data to a base composition signature is useful for certain analyses. As used herein, a "base composition signature" (BCS) is the exact base composition determined from the molecular mass of a bioagent identifying amplicon. In one embodiment, a BCS provides an index of a specific gene
20 in a specific organism.

Base compositions, like sequences, vary slightly from isolate to isolate within species. It is possible to manage this diversity by building "base composition probability clouds" around the composition constraints for each species. This permits identification of organisms in a fashion similar to sequence analysis. A "pseudo four-dimensional plot" can be used to visualize the
25 concept of base composition probability clouds (Figure 18). Optimal primer design requires optimal choice of bioagent identifying amplicons and maximizes the separation between the base composition signatures of individual bioagents. Areas where clouds overlap indicate regions that may result in a misclassification, a problem which is overcome by selecting primers that provide information from different bioagent identifying amplicons, ideally maximizing the separation of
30 base compositions. Thus, one aspect of the utility of an analysis of base composition probability clouds is that it provides a means for screening primer sets in order to avoid potential misclassifications of BCS and bioagent identity. Another aspect of the utility of base composition probability clouds is that they provide a means for predicting the identity of a

bioagent whose exact measured BCS was not previously observed and/or indexed in a BCS database due to evolutionary transitions in its nucleic acid sequence.

It is important to note that, in contrast to probe-based techniques, mass spectrometry determination of base composition does not require prior knowledge of the composition in order to make the measurement, only to interpret the results. In this regard, the present invention provides bioagent classifying information similar to DNA sequencing and phylogenetic analysis at a level sufficient to detect and identify a given bioagent. Furthermore, the process of determination of a previously unknown BCS for a given bioagent (for example, in a case where sequence information is unavailable) has downstream utility by providing additional bioagent indexing information with which to populate BCS databases. The process of future bioagent identification is thus greatly improved as more BCS indexes become available in the BCS databases.

Another embodiment of the present invention is a method of surveying bioagent samples that enables detection and identification of all bacteria for which sequence information is available using a set of twelve broad-range intelligent PCR primers. Six of the twelve primers are "broad range survey primers" herein defined as primers targeted to broad divisions of bacteria (for example, the *Bacillus/Clostridia* group or gamma-proteobacteria). The other six primers of the group of twelve primers are "division-wide" primers herein defined as primers that provide more focused coverage and higher resolution. This method enables identification of nearly 100% of known bacteria at the species level. A further example of this embodiment of the present invention is a method herein designated "survey/drill-down" wherein a subspecies characteristic for detected bioagents is obtained using additional primers. Examples of such a subspecies characteristic include but are not limited to: antibiotic resistance, pathogenicity island, virulence factor, strain type, sub-species type, and clade group. Using the survey/drill-down method, bioagent detection, confirmation and a subspecies characteristic can be provided within hours. Moreover, the survey/drill-down method can be focused to identify bioengineering events such as the insertion of a toxin gene into a bacterial species that does not normally make the toxin.

The present methods allow extremely rapid and accurate detection and identification of bioagents compared to existing methods. Furthermore, this rapid detection and identification is possible even when sample material is impure. The methods leverage ongoing biomedical research in virulence, pathogenicity, drug resistance and genome sequencing into a method which provides greatly improved sensitivity, specificity and reliability compared to existing

methods, with lower rates of false positives. Thus, the methods are useful in a wide variety of fields, including, but not limited to, those fields discussed below.

In other embodiments of the invention, the methods disclosed herein can identify infectious agents in biological samples. At least a first biological sample containing at least a first unidentified infectious agent is obtained. An identification analysis is carried out on the sample, whereby the first infectious agent in the first biological sample is identified. More particularly, a method of identifying an infectious agent in a biological entity is provided. An identification analysis is carried out on a first biological sample obtained from the biological entity, whereby at least one infectious agent in the biological sample from the biological entity is identified. The obtaining and the performing steps are, optionally, repeated on at least one additional biological sample from the biological entity.

The present invention also provides methods of identifying an infectious agent that is potentially the cause of a health condition in a biological entity. An identification analysis is carried out on a first test sample from a first infectious agent differentiating area of the biological entity, whereby at least one infectious agent is identified. The obtaining and the performing steps are, optionally, repeated on an additional infectious agent differentiating area of the biological entity.

Biological samples include, but are not limited to, hair, mucosa, skin, nail, blood, saliva, rectal, lung, stool, urine, breath, nasal, ocular sample, or the like. In some embodiments, one or more biological samples are analyzed by the methods described herein. The biological sample(s) contain at least a first unidentified infectious agent and may contain more than one infectious agent. The biological sample(s) are obtained from a biological entity. The biological sample can be obtained by a variety of manners such as by biopsy, swabbing, and the like. The biological samples may be obtained by a physician in a hospital or other health care environment. The physician may then perform the identification analysis or send the biological sample to a laboratory to carry out the analysis.

Biological entities include, but are not limited to, a mammal, a bird, or a reptile. The biological entity may be a cow, horse, dog, cat, or a primate. The biological entity can also be a human. The biological entity may be living or dead.

An infectious agent differentiating area is any area or location within a biological entity that can distinguish between a harmful versus normal health condition. An infectious agent differentiating area can be a region or area of the biological entity whereby an infectious agent is more likely to predominate from another region or area of the biological entity. For example, infectious agent differentiating areas may include the blood vessels of the heart (heart disease,

coronary artery disease, etc.), particular portions of the digestive system (ulcers, Crohn's disease, etc.), liver (hepatitis infections), and the like. In some embodiments, one or more biological samples from a plurality of infectious agent differentiating areas is analyzed the methods described herein.

5 Infectious agents of the invention may potentially cause a health condition in a biological entity. Health conditions include any condition, syndrome, illness, disease, or the like, identified currently or in the future by medical personnel. Infectious agents include, but are not limited to, bacteria, viruses, parasites, fungi, and the like.

In other embodiments of the invention, the methods disclosed herein can be used to
10 screen blood and other bodily fluids and tissues for pathogenic and non-pathogenic bacteria, viruses, parasites, fungi and the like. Animal samples, including but not limited to, blood and other bodily fluid and tissue samples, can be obtained from living animals, who are either known or not known to or suspected of having a disease, infection, or condition. Alternately, animal samples such as blood and other bodily fluid and tissue samples can be obtained from deceased
15 animals. Blood samples can be further separated into plasma or cellular fractions and further screened as desired. Bodily fluids and tissues can be obtained from any part of the animal or human body. Animal samples can be obtained from, for example, mammals and humans.

Clinical samples are analyzed for disease causing bioagents and biowarfare pathogens simultaneously with detection of bioagents at levels as low as 100-1000 genomic copies in
20 complex backgrounds with throughput of approximately 100-300 samples with simultaneous detection of bacteria and viruses. Such analyses provide additional value in probing bioagent genomes for unanticipated modifications. These analyses are carried out in reference labs, hospitals and the LRN laboratories of the public health system in a coordinated fashion, with the ability to report the results via a computer network to a common data-monitoring center in real
25 time. Clonal propagation of specific infectious agents, as occurs in the epidemic outbreak of infectious disease, can be tracked with base composition signatures, analogous to the pulse field gel electrophoresis fingerprinting patterns used in tracking the spread of specific food pathogens in the Pulse Net system of the CDC (Swaminathan et al., Emerging Infectious Diseases, 2001, 7, 382-389). The present invention provides a digital barcode in the form of a series of base
30 composition signatures, the combination of which is unique for each known organism. This capability enables real-time infectious disease monitoring across broad geographic locations, which may be essential in a simultaneous outbreak or attack in different cities.

In other embodiments of the invention, the methods disclosed herein can be used for detecting the presence of pathogenic and non-pathogenic bacteria, viruses, parasites, fungi and

the like in organ donors and/or in organs from donors. Such examination can result in the prevention of the transfer of, for example, viruses such as West Nile virus, hepatitis viruses, human immunodeficiency virus, and the like from a donor to a recipient via a transplanted organ. The methods disclosed herein can also be used for detection of host versus graft or graft versus
5 host rejection issues related to organ donors by detecting the presence of particular antigens in either the graft or host known or suspected of causing such rejection. In particular, the bioagents in this regard are the antigens of the major histocompatibility complex, such as the HLA antigens. The present methods can also be used to detect and track emerging infectious diseases, such as West Nile virus infection, HIV-related diseases.

10 In other embodiments of the invention, the methods disclosed herein can be used for pharmacogenetic analysis and medical diagnosis including, but not limited to, cancer diagnosis based on mutations and polymorphisms, drug resistance and susceptibility testing, screening for and/or diagnosis of genetic diseases and conditions, and diagnosis of infectious diseases and conditions. In context of the present invention, pharmacogenetics is defined as the study of
15 variability in drug response due to genetic factors. Pharmacogenetic investigations are often based on correlating patient outcome with variations in genes involved in the mode of action of a given drug. For example, receptor genes, or genes involved in metabolic pathways. The methods of the present invention provide a means to analyze the DNA of a patient to provide the basis for pharmacogenetic analysis.

20 The present method can also be used to detect single nucleotide polymorphisms (SNPs), or multiple nucleotide polymorphisms, rapidly and accurately. A SNP is defined as a single base pair site in the genome that is different from one individual to another. The difference can be expressed either as a deletion, an insertion or a substitution, and is frequently linked to a disease state. Because they occur every 100-1000 base pairs, SNPs are the most frequently found type of
25 genetic marker in the human genome.

For example, sickle cell anemia results from an A-T transition, which encodes a valine rather than a glutamic acid residue. Oligonucleotide primers may be designed such that they bind to sequences that flank a SNP site, followed by nucleotide amplification and mass determination of the amplified product. Because the molecular masses of the resulting product from an
30 individual who does not have sickle cell anemia is different from that of the product from an individual who has the disease, the method can be used to distinguish the two individuals. Thus, the method can be used to detect any known SNP in an individual and thus diagnose or determine increased susceptibility to a disease or condition.

In one embodiment, blood is drawn from an individual and peripheral blood mononuclear cells (PBMC) are isolated and simultaneously tested, such as in a high-throughput screening method, for one or more SNPs using appropriate primers based on the known sequences which flank the SNP region. The National Center for Biotechnology Information
5 maintains a publicly available database of SNPs on the world wide web of the Internet at, for example, "ncbi.nlm.nih.gov/SNP/."

The method of the present invention can also be used for blood typing. The gene encoding A, B or O blood type can differ by four single nucleotide polymorphisms. If the gene contains the sequence CGTGGTGACCCTT (SEQ ID NO:5), antigen A results. If the gene
10 contains the sequence CGTCGTCACCGCTA (SEQ ID NO:6) antigen B results. If the gene contains the sequence CGTGGT-ACCCCTT (SEQ ID NO:7), blood group O results ("-" indicates a deletion). These sequences can be distinguished by designing a single primer pair which flanks these regions, followed by amplification and mass determination.

The method of the present invention can also be used for detection and identification of
15 blood-borne pathogens such as *Staphylococcus aureus* for example.

The method of the present invention can also be used for strain typing of respiratory pathogens in epidemic surveillance. Group A streptococci (GAS), or *Streptococcus pyogenes*, is one of the most consequential causes of respiratory infections because of prevalence and ability to cause disease with complications such as acute rheumatic fever and acute glomerulonephritis. GAS
20 also causes infections of the skin (*impetigo*) and, in rare cases, invasive disease such as necrotizing fasciitis and toxic shock syndrome. Despite many decades of study, the underlying microbial ecology and natural selection that favors enhanced virulence and explosive GAS outbreaks is still poorly understood. The ability to detect GAS and multiple other pathogenic and non-pathogenic bacteria and viruses in patient samples would greatly facilitate our
25 understanding of GAS epidemics. It is also essential to be able to follow the spread of virulent strains of GAS in populations and to distinguish virulent strains from less virulent or avirulent streptococci that colonize the nose and throat of asymptomatic individuals at a frequency ranging from 5-20% of the population (Bisno, A. L. (1995) in Principles and Practice of Infectious Diseases, eds. Mandell, G. L., Bennett, J. E. & Dolin, R. (Churchill Livingstone, New York), Vol.
30 2, pp. 1786-1799). Molecular methods have been developed to type GAS based upon the sequence of the emm gene that encodes the M-protein virulence factor (Beall et al., J. Clin. Micro., 1996, 34, 953-958; Beall et al., J. Clin. Micro., 1997, 35, 1231-1235; and Facklam et al., Emerging Infectious Diseases, 1999, 5, 247-253). Using this molecular classification, over 150 different emm-types are defined and correlated with phenotypic properties of thousands of GAS

isolates (www.cdc.gov/ncidod/biotech/strep/strepindex.html) (Facklam et al., Clinical Infectious Diseases, 2002, 34, 28-38). Recently, a strategy known as Multi Locus Sequence Typing (MLST) was developed to follow the molecular Epidemiology of GAS. In MLST, internal fragments of seven housekeeping genes are amplified, sequenced, and compared to a database of previously studied isolates (www.test.mlst.net/).

The present invention enables an emm-typing process to be carried out directly from throat swabs for a large number of samples within 12 hours, allowing strain tracking of an ongoing epidemic, even if geographically dispersed, on a larger scale than ever before achievable.

10 In another embodiment, the present invention can be employed in the serotyping of viruses including, but not limited to, adenoviruses. Adenoviruses are DNA viruses that cause over 50% of febrile respiratory illnesses in military recruits. Human adenoviruses are divided into six major serogroups (A through F), each containing multiple strain types. Despite the prevalence of adenoviruses, there are no rapid methods for detecting and serotyping
15 adenoviruses.

In another embodiment, the present invention can be employed in distinguishing between members of the *Orthopoxvirus* genus. Smallpox is caused by the *Variola* virus. Other members of the genus include *Vaccinia*, *Monkeypox*, *Camelpox*, and *Cowpox*. All are capable of infecting humans, thus, a method capable of identifying and distinguishing among members of
20 the *Orthopox* genus is a worthwhile objective.

In another embodiment, the present invention can be employed in distinguishing between viral agents of viral hemorrhagic fevers (VHF). VHF agents include, but are not limited to, *Filoviridae* (Marburg virus and Ebola virus), *Arenaviridae* (Lassa, Junin, Machupo, Sabia, and Guanarito viruses), *Bunyaviridae* (Crimean-Congo hemorrhagic fever virus (CCHFV), Rift
25 Valley fever virus, and Hanta viruses), and *Flaviviridae* (yellow fever virus and dengue virus). Infections by VHF viruses are associated with a wide spectrum of clinical manifestations such as diarrhea, myalgia, cough, headache, pneumonia, encephalopathy, and hepatitis. Filoviruses, arenaviruses, and CCHFV are of particular relevance because they can be transmitted from human to human, thus causing epidemics with high mortality rates (Khan et al., Am. J. Trop.
30 Med. Hyg., 1997, 57, 519-525). In the absence of bleeding or organ manifestation, VHF is clinically difficult to diagnose, and the various etiologic agents can hardly be distinguished by clinical tests. Current approaches to PCR detection of these agents are time-consuming, as they include a separate cDNA synthesis step prior to PCR, agarose gel analysis of PCR products, and in some instances a second round of nested amplification or Southern hybridization. PCRs for

different pathogens have to be run assay by assay due to differences in cycling conditions, which complicate broad-range testing in a short period. Moreover, post-PCR processing or nested PCR steps included in currently used assays increase the risk of false positive results due to carryover contamination (Kwok et al., Nature, 1989, 339, 237-238).

- 5 In another embodiment, the present invention, can be employed in the diagnosis of a plurality of etiologic agents of a disease. An "etiologic agent" is herein defined as a pathogen acting as the causative agent of a disease. Diseases may be caused by a plurality of etiologic agents. For example, recent studies have implicated both human herpesvirus 6 (HHV-6) and the obligate intracellular bacterium *Chlamydia pneumoniae* in the etiology of multiple sclerosis
- 10 (Swanborg, Microbes and Infection, 2002, 4, 1327-1333). The present invention can be applied to the identification of multiple etiologic agents of a disease by, for example, the use of broad range bacterial intelligent primers and division-wide primers (if necessary) for the identification of bacteria such as *Chlamydia pneumoniae* followed by primers directed to viral housekeeping genes for the identification of viruses such as HHV-6, for example.
- 15 In other embodiments of the invention, the methods disclosed herein can be used for detection and identification of pathogens in livestock. Livestock includes, but is not limited to, cows, pigs, sheep, chickens, turkeys, goats, horses and other farm animals. For example, conditions classified by the California Department of Food and Agriculture as emergency conditions in livestock (www.cdffa.ca.gov/ahfss/ah/pdfs/CA_reportable_disease_list_05292002.pdf) include, but are not limited to: Anthrax (*Bacillus anthracis*), Screwworm myiasis (*Cochliomyia hominivorax* or *Chrysomya bezziana*), African trypanosomiasis (Tsetse fly diseases), Bovine babesiosis (*piroplasmosis*), Bovine spongiform encephalopathy (Mad Cow), Contagious bovine pleuropneumonia (*Mycoplasma mycoides mycoides* small colony), Foot-and-mouth disease (Hoof-and-mouth), Heartwater (*Cowdria ruminantium*), Hemorrhagic septicemia
- 25 (*Pasteurella multocida* serotypes B:2 or E:2), Lumpy skin disease, Malignant catarrhal fever (African type), Rift Valley fever, Rinderpest (Cattle plague), Theileriosis (Corridor disease, East Coast fever), Vesicular stomatitis, Contagious agalactia (*Mycoplasma* species), Contagious caprine pleuropneumonia (*Mycoplasma capricolum capripneumoniae*), Nairobi sheep disease, Peste des petits ruminants (Goat plague), Pulmonary adenomatosis (Viral neoplastic pneumonia),
- 30 *Salmonella abortus ovis*, Sheep and goat pox, African swine fever, Classical swine fever (Hog cholera), Japanese encephalitis, Nipah virus, Swine vesicular disease, Teschen disease (*Enterovirus encephalomyelitis*), Vesicular exanthema, Exotic Newcastle disease (Viscerotropic velogenic Newcastle disease), Highly pathogenic avian influenza (Fowl plague), African horse sickness, Dourine (*Trypanosoma equiperdum*), Epizootic lymphangitis (equine blastomycosis,

equine histoplasmosis), Equine piroplasmiasis (*Babesia equi*, *B. caballi*), Glanders (Farcy) (*Pseudomonas mallei*), Hendra virus (Equine morbillivirus), Horse pox, Surra (*Trypanosoma evansi*), Venezuelan equine encephalomyelitis, West Nile Virus, Chronic wasting disease in cervids, and Viral hemorrhagic disease of rabbits (calicivirus)

- 5 Conditions classified by the California Department of Food and Agriculture as regulated conditions in livestock include, but are not limited to: rabies, Bovine brucellosis (*Brucella abortus*), Bovine tuberculosis (*Mycobacterium bovis*), Cattle scabies (multiple types), Trichomonosis (*Trichomonas fetus*), Caprine and ovine brucellosis (excluding *Brucella ovis*), Scrapie, Sheep scabies (Body mange) (*Psoroptes ovis*), Porcine brucellosis (*Brucella suis*),
- 10 Pseudorabies (Aujeszky's disease), Ornithosis (*Psittacosis* or avian *chlamydiosis*) (*Chlamydia psittaci*), Pullorum disease (Fowl typhoid) (*Salmonella gallinarum* and *pullorum*), Contagious equine metritis (*Taylorella equigenitalis*), Equine encephalomyelitis (Eastern and Western equine encephalitis), Equine infectious anemia (Swamp fever), Duck viral enteritis (Duck plague), and Tuberculosis in cervids.
- 15 Additional conditions monitored by the California Department of Food and Agriculture include, but are not limited to: Avian tuberculosis (*Mycobacterium avium*), Echinococcosis/Hydatidosis (*Echinococcus* species), Leptospirosis, Anaplasmosis (*Anaplasma marginale* or *A. centrale*), Bluetongue, Bovine cysticercosis (*Taenia saginata* in humans), Bovine genital campylobacteriosis (*Campylobacter fetus venerealis*), Dermatophilosis
- 20 (Streptothricosis, mycotic dermatitis) (*Dermatophilus congolensis*), Enzootic bovine leukosis (Bovine leukemia virus), Infectious bovine rhinotracheitis (Bovine herpesvirus-1), Johne's disease (Paratuberculosis) (*Mycobacterium avium paratuberculosis*), Malignant catarrhal fever (North American), Q Fever (*Coxiella burnetii*), Caprine (contagious) arthritis/encephalitis, Enzootic abortion of ewes (Ovine chlamydiosis) (*Chlamydia psittaci*), Maedi-Visna (Ovine
- 25 progressive pneumonia), Atrophic rhinitis (*Bordetella bronchiseptica*, *Pasteurella multocida*), Porcine cysticercosis (*Taenia solium* in humans), Porcine reproductive and respiratory syndrome, Transmissible gastroenteritis (coronavirus), Trichinellosis (*Trichinella spiralis*), Avian infectious bronchitis, Avian infectious laryngotracheitis, Duck viral hepatitis, Fowl cholera (*Pasteurella multocida*), Fowl pox, Infectious bursal disease (Gumboro disease), Low
- 30 pathogenic avian influenza, Marek's disease, Mycoplasmosis (*Mycoplasma gallisepticum*), Equine influenza Equine rhinopneumonitis (Equine herpesvirus-1), Equine viral arteritis, and Horse mange (multiple types).

A key problem in determining that an infectious outbreak is the result of a bioterrorist attack is the sheer variety of organisms that might be used by terrorists. According to a recent

review (Taylor et al., Philos. Trans. R. Soc. Lond. B. Biol. Sci., 2001, 356, 983-989), there are over 1400 organisms infectious to humans; most of these have the potential to be used in a deliberate, malicious attack. These numbers do not include numerous strain variants of each organism, bioengineered versions, or pathogens that infect plants or animals. Paradoxically, most of the new technology being developed for detection of biological weapons incorporates a version of quantitative PCR, which is based upon the use of highly specific primers and probes designed to selectively identify specific pathogenic organisms. This approach requires assumptions about the type and strain of bacteria or virus which is expected to be detected. Although this approach will work for the most obvious organisms, like smallpox and anthrax, experience has shown that it is very difficult to anticipate what a terrorist will do.

The present invention can be used to detect and identify any biological agent, including bacteria, viruses, fungi and toxins without prior knowledge of the organism being detected and identified. As one example, where the agent is a biological threat, the information obtained such as the presence of toxin genes, pathogenicity islands and antibiotic resistance genes for example, is used to determine practical information needed for countermeasures. In addition, the methods can be used to identify natural or deliberate engineering events including chromosome fragment swapping, molecular breeding (gene shuffling) and emerging infectious diseases. The present invention provides broad-function technology that may be the only practical means for rapid diagnosis of disease caused by a biowarfare or bioterrorist attack, especially an attack that might otherwise be missed or mistaken for a more common infection.

Bacterial biological warfare agents capable of being detected by the present methods include, but are not limited to, *Bacillus anthracis* (anthrax), *Yersinia pestis* (pneumonic plague), *Francisella tularensis* (tularemia), *Brucella suis*, *Brucella abortus*, *Brucella melitensis* (undulant fever), *Burkholderia mallei* (glanders), *Burkholderia pseudomallei* (melioidosis), *Salmonella typhi* (typhoid fever), *Rickettsia typhi* (epidemic typhus), *Rickettsia prowasekii* (endemic typhus) and *Coxiella burnetii* (Q fever), *Rhodobacter capsulatus*, *Chlamydia pneumoniae*, *Escherichia coli*, *Shigella dysenteriae*, *Shigella flexneri*, *Bacillus cereus*, *Clostridium botulinum*, *Coxiella burnetii*, *Pseudomonas aeruginosa*, *Legionella pneumophila*, and *Vibrio cholerae*.

Besides 16S and 23S rRNA, other target regions suitable for use in the present invention for detection of bacteria include, but are not limited to, 5S rRNA and RNase P (Figure 3).

Fungal biowarfare agents include, but are not limited to, *Coccidioides immitis* (Coccidioidomycosis), and *Magnaporthe grisea*.

Biological warfare toxin genes capable of being detected by the methods of the present invention include, but are not limited to, botulinum toxin, T-2 mycotoxins, ricin, staph enterotoxin B, shigatoxin, abrin, aflatoxin, *Clostridium perfringens* epsilon toxin, conotoxins, diacetoxyscirpenol, tetrodotoxin and saxitoxin.

- 5 Parasites that could be used in biological warfare include, but are not limited to: *Ascaris suum*, *Giardia lamblia*, *Cryptosporidium*, and *Schistosoma*.

Biological warfare viral threat agents are mostly RNA viruses (positive-strand and negative-strand), with the exception of smallpox. Every RNA virus is a family of related viruses (quasispecies). These viruses mutate rapidly and the potential for engineered strains (natural or
10 deliberate) is very high. RNA viruses cluster into families that have conserved RNA structural domains on the viral genome (e.g., virion components, accessory proteins) and conserved housekeeping genes that encode core viral proteins including, for single strand positive strand RNA viruses, RNA-dependent RNA polymerase, double stranded RNA helicase, chymotrypsin-like and papain-like proteases and methyltransferases. "Housekeeping genes" refers to genes that
15 are generally always expressed and thought to be involved in routine cellular metabolism.

Examples of (-)-strand RNA viruses include, but are not limited to, arenaviruses (e.g., sabia virus, lassa fever, Machupo, Argentine hemorrhagic fever, flexal virus), bunyaviruses (e.g., hantavirus, nairovirus, phlebovirus, hantaan virus, Congo-crimean hemorrhagic fever, rift valley fever), and mononegavirales (e.g., filovirus, paramyxovirus, ebola virus, Marburg, equine
20 morbillivirus).

Examples of (+)-strand RNA viruses include, but are not limited to, picornaviruses (e.g., coxsackievirus, echovirus, human coxsackievirus A, human echovirus, human enterovirus, human poliovirus, hepatitis A virus, human parechovirus, human rhinovirus), astroviruses (e.g., human astrovirus), calciviruses (e.g., chiba virus, chitta virus, human calcivirus, norwalk virus),
25 nidovirales (e.g., human coronavirus, human torovirus), flaviviruses (e.g., dengue virus 1-4, Japanese encephalitis virus, Kyasanur forest disease virus, Murray Valley encephalitis virus, Rocio virus, St. Louis encephalitis virus, West Nile virus, yellow fever virus, hepatitis c virus) and togaviruses (e.g., Chikugunya virus, Eastern equine encephalitis virus, Mayaro virus, O'nyong-nyong virus, Ross River virus, Venezuelan equine encephalitis virus, Rubella virus,
30 hepatitis E virus). The hepatitis C virus has a 5'-untranslated region of 340 nucleotides, an open reading frame encoding 9 proteins having 3010 amino acids and a 3'-untranslated region of 240 nucleotides. The 5'-UTR and 3'-UTR are 99% conserved in hepatitis C viruses.

In one embodiment, the target gene is an RNA-dependent RNA polymerase or a helicase encoded by (+)-strand RNA viruses, or RNA polymerase from a (-)-strand RNA virus.

(+)-strand RNA viruses are double stranded RNA and replicate by RNA-directed RNA synthesis using RNA-dependent RNA polymerase and the positive strand as a template. Helicase unwinds the RNA duplex to allow replication of the single stranded RNA. These viruses include viruses from the family picornaviridae (e.g., poliovirus, coxsackievirus, echovirus), togaviridae (e.g., alphavirus, flavivirus, rubivirus), arenaviridae (e.g., lymphocytic choriomeningitis virus, lassa fever virus), coronaviridae (e.g., human respiratory virus) and Hepatitis A virus. The genes encoding these proteins comprise variable and highly conserved regions that flank the variable regions.

In one embodiment, the method can be used to detect the presence of antibiotic resistance and/or toxin genes in a bacterial species. For example, *Bacillus anthracis* comprising a tetracycline resistance plasmid and plasmids encoding one or both anthracis toxins (px01 and/or px02) can be detected by using antibiotic resistance primer sets and toxin gene primer sets. If the *B. anthracis* is positive for tetracycline resistance, then a different antibiotic, for example quinalone, is used.

While the present invention has been described with specificity in accordance with certain of its embodiments, the following examples serve only to illustrate the invention and are not intended to limit the same.

EXAMPLES

Example 1: Nucleic Acid Isolation and PCR

In one embodiment, nucleic acid is isolated from the organisms and amplified by PCR using standard methods prior to BCS determination by mass spectrometry. Nucleic acid is isolated, for example, by detergent lysis of bacterial cells, centrifugation and ethanol precipitation. Nucleic acid isolation methods are described in, for example, *Current Protocols in Molecular Biology* (Ausubel et al.) and *Molecular Cloning: A Laboratory Manual* (Sambrook et al.). The nucleic acid is then amplified using standard methodology, such as PCR, with primers which bind to conserved regions of the nucleic acid which contain an intervening variable sequence as described below.

General Genomic DNA Sample Prep Protocol: Raw samples are filtered using Supor-200 0.2 μ m membrane syringe filters (VWR International). Samples are transferred to 1.5 ml eppendorf tubes pre-filled with 0.45 g of 0.7 mm Zirconia beads followed by the addition of 350 μ l of ATL buffer (Qiagen, Valencia, CA). The samples are subjected to bead beating for 10 minutes at a frequency of 19 1/s in a Retsch Vibration Mill (Retsch). After centrifugation,

samples are transferred to an S-block plate (Qiagen) and DNA isolation is completed with a BioRobot 8000 nucleic acid isolation robot (Qiagen).

Swab Sample Protocol: Allegiance S/P brand culture swabs and collection/transport system are used to collect samples. After drying, swabs are placed in 17x100 mm culture tubes
5 (VWR International) and the genomic nucleic acid isolation is carried out automatically with a Qiagen Mdx robot and the Qiagen QIAamp DNA Blood BioRobot Mdx genomic preparation kit (Qiagen, Valencia, CA).

Example 2: Mass spectrometry

10 *FTICR Instrumentation:* The FTICR instrument is based on a 7 tesla actively shielded superconducting magnet and modified Bruker Daltonics Apex II 70e ion optics and vacuum chamber. The spectrometer is interfaced to a LEAP PAL autosampler and a custom fluidics control system for high throughput screening applications. Samples are analyzed directly from 96-well or 384-well microtiter plates at a rate of about 1 sample/minute. The Bruker data-
15 acquisition platform is supplemented with a lab-built ancillary NT datastation which controls the autosampler and contains an arbitrary waveform generator capable of generating complex rf-excite waveforms (frequency sweeps, filtered noise, stored waveform inverse Fourier transform (SWIFT), etc.) for sophisticated tandem MS experiments. For oligonucleotides in the 20-30-mer regime typical performance characteristics include mass resolving power in excess of 100,000
20 (FWHM), low ppm mass measurement errors, and an operable m/z range between 50 and 5000 m/z .

Modified ESI Source: In sample-limited analyses, analyte solutions are delivered at 150 nL/minute to a 30 mm i.d. fused-silica ESI emitter mounted on a 3-D micromanipulator. The ESI ion optics consists of a heated metal capillary, an rf-only hexapole, a skimmer cone, and an
25 auxiliary gate electrode. The 6.2 cm rf-only hexapole is comprised of 1 mm diameter rods and is operated at a voltage of 380 Vpp at a frequency of 5 MHz. A lab-built electro-mechanical shutter can be employed to prevent the electrospray plume from entering the inlet capillary unless triggered to the "open" position via a TTL pulse from the data station. When in the "closed" position, a stable electrospray plume is maintained between the ESI emitter and the face of the
30 shutter. The back face of the shutter arm contains an elastomeric seal that can be positioned to form a vacuum seal with the inlet capillary. When the seal is removed, a 1 mm gap between the shutter blade and the capillary inlet allows constant pressure in the external ion reservoir regardless of whether the shutter is in the open or closed position. When the shutter is triggered, a "time slice" of ions is allowed to enter the inlet capillary and is subsequently accumulated in

the external ion reservoir. The rapid response time of the ion shutter (< 25 ms) provides reproducible, user defined intervals during which ions can be injected into and accumulated in the external ion reservoir.

Apparatus for Infrared Multiphoton Dissociation: A 25 watt CW CO₂ laser operating at 5 10.6 μm has been interfaced to the spectrometer to enable infrared multiphoton dissociation (IRMPD) for oligonucleotide sequencing and other tandem MS applications. An aluminum optical bench is positioned approximately 1.5 m from the actively shielded superconducting magnet such that the laser beam is aligned with the central axis of the magnet. Using standard IR-compatible mirrors and kinematic mirror mounts, the unfocused 3 mm laser beam is aligned 10 to traverse directly through the 3.5 mm holes in the trapping electrodes of the FTICR trapped ion cell and longitudinally traverse the hexapole region of the external ion guide finally impinging on the skimmer cone. This scheme allows IRMPD to be conducted in an m/z selective manner in the trapped ion cell (e.g. following a SWIFT isolation of the species of interest), or in a broadband mode in the high pressure region of the external ion reservoir where collisions with 15 neutral molecules stabilize IRMPD-generated metastable fragment ions resulting in increased fragment ion yield and sequence coverage.

Example 3: Identification of Bioagents

Table 2 shows a small cross section of a database of calculated molecular masses for 20 over 9 primer sets and approximately 30 organisms. The primer sets were derived from rRNA alignment. Examples of regions from rRNA consensus alignments are shown in Figures 1A-1C. Lines with arrows are examples of regions to which intelligent primer pairs for PCR are designed. The primer pairs are >95% conserved in the bacterial sequence database (currently over 10,000 organisms). The intervening regions are variable in length and/or composition, thus 25 providing the base composition "signature" (BCS) for each organism. Primer pairs were chosen so the total length of the amplified region is less than about 80-90 nucleotides. The label for each primer pair represents the starting and ending base number of the amplified region on the consensus diagram.

Included in the short bacterial database cross-section in Table 2 are many well known 30 pathogens/biowarfare agents (shown in bold/red typeface) such as *Bacillus anthracis* or *Yersinia pestis* as well as some of the bacterial organisms found commonly in the natural environment such as *Streptomyces*. Even closely related organisms can be distinguished from each other by the appropriate choice of primers. For instance, two low G+C organisms, *Bacillus anthracis* and

Staph aureus, can be distinguished from each other by using the primer pair defined by 16S_1337 or 23S_855 (ΔM of 4 Da).

Table 2: Cross Section Of A Database Of Calculated Molecular Masses¹

Primer Regions → Bug Name	18S_971	6S_1100	6S_1337	6S_1294	6S_1228	23S_1021	23S_855	23S_193	23S_115
<i>Acinetobacter calcoaceticus</i>	55619.1	55004	28446.7	35854.9	51295.4	30299	42654	39557.5	54999
<i>Bacillus anthracis</i>	55095	54388	28448	35238	51296	30295	42651	39560	56850
<i>Bacillus cereus</i>	55622.1	54387.9	28447.8	35854.9	51295.4	30295	42651	39560.5	56850.3
<i>Bordetella bronchiseptica</i>	56857.3	51300.4	28448.7	35857.9	51307.4	30299	42653	39559.5	51820.6
<i>Borrelia burgdorferi</i>	56231.2	55621.1	28440.7	35852.9	51295.4	30297	42029.9	38941.4	52523.6
<i>Brucella abortus</i>	58098	55011	28448	35854	50683				
<i>Campylobacter jejuni</i>	58088.6	54386.9	28061.8	35856.9	50674.3	30294	42032.9	39558.5	45732.5
<i>Chlamydia pneumoniae</i>	55000	55007	29063	35855	50676	30295	42036	38941	56230
<i>Clostridium botulinum</i>	55006	53767	28445	35855	51281	30300	42656	39562	54999
<i>Clostridium difficile</i>	56855.3	54386.9	28444.7	35853.9	51295.4	30294	41417.8	39558.5	56812.2
<i>Enterococcus faecalis</i>	55620.1	54387.9	28447.6	35858.9	51295.4	30297	42652	39559.5	56849.3
<i>Escherichia coli</i>	55622	55009	28445	35857	51301	30301	42656	39562	54999
<i>Francisella tularensis</i>	53769	54385	28445	35856	51298				
<i>Haemophilus influenzae</i>	55620.1	55006	28444.7	35855.9	51298.4	30298	42656	39560.5	55613.1
<i>Klebsiella pneumoniae</i>	55622.1	55008	28442.7	35856.9	51297.4	30300	42655	39562.5	55000
<i>Legionella pneumophila</i>	55618	55626	28446	35857	51303				
<i>Mycobacterium avium</i>	54390.9	55631.1	29064.8	35858.9	51815.5	30298	42656	38942.4	56241.2
<i>Mycobacterium leprae</i>	54389.9	55629.1	29064.8	35860.9	51917.5	30298	42656	39559.5	56240.2
<i>Mycobacterium tuberculosis</i>	54390.9	55629.1	29064.8	35860.9	51301.4	30299	42658	39560.5	56243.2
<i>Mycoplasma genitalium</i>	53143.7	45115.4	29081.8	35854.9	50871.3	30294	43264.1	39558.5	56842.4
<i>Mycoplasma pneumoniae</i>	53143.7	45118.4	29081.8	35854.9	50673.3	30294	43264.1	39559.5	56843.4
<i>Neisseria gonorrhoeae</i>	55627.1	54389.9	28445.7	35855.9	51302.4	30300	42849	39561.5	55000
<i>Pseudomonas aeruginosa</i>	55623	55010	28443	35858	51301	30298	43272	39558	55619
<i>Rickettsia prowazekii</i>	58093	55621	28448	35853	50677	30293	42650	39559	53139
<i>Rickettsia rickettsii</i>	58094	55623	28448	35853	50679	30293	42648	39559	53755
<i>Salmonella typhimurium</i>	55622	55005	28445	35857	51301	30301	42658		
<i>Shigella dysenteriae</i>	55623	55009	28444	35857	51301				
<i>Staphylococcus aureus</i>	56854.3	54386.9	28443.7	35852.9	51294.4	30298	42655	39559.5	57466.4
<i>Streptomyces</i>	54389.9	58341.6	29063.8	35858.9	51300.4			39563.5	58864.3
<i>Treponema pallidum</i>	56245.2	55631.1	28445.7	35851.9	51297.4	30299	42034.9	38939.4	57473.4
<i>Vibrio cholerae</i>	55625	55626	28443	35857	52536	29063	30303	35241	50675
<i>Vibrio parahaemolyticus</i>	54384.9	55626.1	28444.7	34620.7	50084.2				
<i>Yersinia pestis</i>	55620	55626	28443	35857	51299				

- 5 ¹ Molecular mass distribution of PCR amplified regions for a selection of organisms (rows) across various primer pairs (columns). Pathogens are shown in bold. Empty cells indicate presently incomplete or missing data.

Figure 6 shows the use of ESI-FT-ICR MS for measurement of exact mass. The spectra from 46mer PCR products originating at position 1337 of the 16S rRNA from *S. aureus* (upper) and *B. anthracis* (lower) are shown. These data are from the region of the spectrum containing signals from the $[M-8H]^8+$ charge states of the respective 5'-3' strands. The two strands differ by two (AT→CG) substitutions, and have measured masses of 14206.396 and 14208.373 + 0.010 Da, respectively. The possible base compositions derived from the masses of the forward and reverse strands for the *B. anthracis* products are listed in Table 3.

15 **Table 3: Possible base composition for *B. anthracis* products**

Calc. Mass	Error	Base Comp.
14208.2935	0.079520	A1 G17 C10 T18
14208.3160	0.056980	A1 G20 C15 T10

14208.3386	0.034440	A1 G23 C20 T2
14208.3074	0.065560	A6 G11 C3 T26
14208.3300	0.043020	A6 G14 C8 T18
14208.3525	0.020480	A6 G17 C13 T10
14208.3751	0.002060	A6 G20 C18 T2
14208.3439	0.029060	A11 G8 C1 T26
14208.3665	0.006520	A11 G11 C6 T18
14208.3890	0.016020	A11 G14 C11 T10
14208.4116	0.038560	A11 G17 C16 T2
14208.4030	0.029980	A16 G8 C4 T18
14208.4255	0.052520	A16 G11 C9 T10
14208.4481	0.075060	A16 G14 C14 T2
14208.4395	0.066480	A21 G5 C2 T18
14208.4620	0.089020	A21 G8 C7 T10
14079.2624	0.080600	A0 G14 C13 T19
14079.2849	0.058060	A0 G17 C18 T11
14079.3075	0.035520	A0 G20 C23 T3
14079.2538	0.089180	A5 G5 C1 T35
14079.2764	0.066640	A5 G8 C6 T27
14079.2989	0.044100	A5 G11 C11 T19
14079.3214	0.021560	A5 G14 C16 T11
14079.3440	0.000980	A5 G17 C21 T3
14079.3129	0.030140	A10 G5 C4 T27
14079.3354	0.007600	A10 G8 C9 T19
14079.3579	0.014940	A10 G11 C14 T11
14079.3805	0.037480	A10 G14 C19 T3
14079.3494	0.006360	A15 G2 C2 T27
14079.3719	0.028900	A15 G5 C7 T19
14079.3944	0.051440	A15 G8 C12 T11
14079.4170	0.073980	A15 G11 C17 T3
14079.4084	0.065400	A20 G2 C5 T19
14079.4309	0.087940	A20 G5 C10 T13

Among the 16 compositions for the forward strand and the 18 compositions for the reverse strand that were calculated, only one pair (shown in **bold**) are complementary, corresponding to the actual base compositions of the *B. anthracis* PCR products.

5 Example 4: BCS of Region from *Bacillus anthracis* and *Bacillus cereus*

A conserved *Bacillus* region from *B. anthracis* (A₁₄G₉C₁₄T₉) and *B. cereus* (A₁₅G₉C₁₃T₉) having a C to A base change was synthesized and subjected to ESI-TOF MS. The results are shown in Figure 7 in which the two regions are clearly distinguished using the method of the present invention (MW=14072.26 vs. 14096.29).

10

Example 5: Identification of additional bioagents

In other examples of the present invention, the pathogen *Vibrio cholera* can be distinguished from *Vibrio parahaemolyticus* with $\Delta M > 600$ Da using one of three 16S primer sets shown in Table 2 (16S_971, 16S_1228 or 16S_1294) as shown in Table 4. The two mycoplasma
15 species in the list (*M. genitalium* and *M. pneumoniae*) can also be distinguished from each other, as can the three mycobacteriae. While the direct mass measurements of amplified products can identify and distinguish a large number of organisms, measurement of the base composition signature provides dramatically enhanced resolving power for closely related organisms. In cases such as *Bacillus anthracis* and *Bacillus cereus* that are virtually indistinguishable from each
20 other based solely on mass differences, compositional analysis or fragmentation patterns are used to resolve the differences. The single base difference between the two organisms yields different fragmentation patterns, and despite the presence of the ambiguous/unidentified base N at position 20 in *B. anthracis*, the two organisms can be identified.

Tables 4a-b show examples of primer pairs from Table 1 which distinguish pathogens
25 from background.

Table 4a

Organism name	23S_855	16S_1337	23S_1021
<i>Bacillus anthracis</i>	42650.98	28447.65	30294.98
<i>Staphylococcus aureus</i>	42654.97	28443.67	30297.96

Table 4b

Organism name	16S_971	16S_1294	16S_1228
<i>Vibrio cholerae</i>	55625.09	35856.87	52535.59
<i>Vibrio parahaemolyticus</i>	54384.91	34620.67	50064.19

Table 5 shows the expected molecular weight and base composition of region 16S_1100-1188 in *Mycobacterium avium* and *Streptomyces sp.*

5

Table 5

Region	Organism name	Length	Molecular weight	Base comp.
16S_1100-1188	<i>Mycobacterium avium</i>	82	25624.1728	A ₁₆ G ₃₂ C ₁₈ T ₁₆
16S_1100-1188	<i>Streptomyces sp.</i>	96	29904.871	A ₁₇ G ₃₈ C ₂₇ T ₁₄

Table 6 shows base composition (single strand) results for 16S_1100-1188 primer amplification reactions different species of bacteria. Species which are repeated in the table 10 (e.g., *Clostridium botulinum*) are different strains which have different base compositions in the 16S_1100-1188 region.

Table 6

Organism name	Base comp.	Organism name	Base comp.
<i>Mycobacterium avium</i>	A ₁₆ G ₃₂ C ₁₈ T ₁₆	<i>Vibrio cholerae</i>	A ₂₃ G ₃₀ C ₂₁ T ₁₆
<i>Streptomyces sp.</i>	A ₁₇ G ₃₈ C ₂₇ T ₁₄	<i>Aeromonas hydrophila</i>	A ₂₃ G ₃₁ C ₂₁ T ₁₅
<i>Ureaplasma urealyticum</i>	A ₁₈ G ₃₀ C ₁₇ T ₁₇	<i>Aeromonas salmonicida</i>	A ₂₃ G ₃₁ C ₂₁ T ₁₅
<i>Streptomyces sp.</i>	A ₁₉ G ₃₆ C ₂₄ T ₁₈	<i>Mycoplasma genitalium</i>	A ₂₄ G ₁₉ C ₁₂ T ₁₈
<i>Mycobacterium leprae</i>	A ₂₀ G ₃₂ C ₂₂ T ₁₆	<i>Clostridium botulinum</i>	A ₂₄ G ₂₅ C ₁₈ T ₂₀
<i>M. tuberculosis</i>	A ₂₀ G ₃₃ C ₂₁ T ₁₆	<i>Bordetella bronchiseptica</i>	A ₂₄ G ₂₆ C ₁₉ T ₁₄
<i>Nocardia asteroides</i>	A ₂₀ G ₃₃ C ₂₁ T ₁₆	<i>Francisella tularensis</i>	A ₂₄ G ₂₆ C ₁₉ T ₁₉
<i>Fusobacterium necroforum</i>	A ₂₁ G ₂₆ C ₂₂ T ₁₈	<i>Bacillus anthracis</i>	A ₂₄ G ₂₆ C ₂₀ T ₁₈
<i>Listeria monocytogenes</i>	A ₂₁ G ₂₇ C ₁₉ T ₁₉	<i>Campylobacter jejuni</i>	A ₂₄ G ₂₆ C ₂₀ T ₁₈
<i>Clostridium botulinum</i>	A ₂₁ G ₂₇ C ₁₉ T ₂₁	<i>Staphylococcus aureus</i>	A ₂₄ G ₂₆ C ₂₀ T ₁₈
<i>Neisseria gonorrhoeae</i>	A ₂₁ G ₂₈ C ₂₁ T ₁₈	<i>Helicobacter pylori</i>	A ₂₄ G ₂₆ C ₂₀ T ₁₉
<i>Bartonella quintana</i>	A ₂₁ G ₃₀ C ₂₂ T ₁₆	<i>Helicobacter pylori</i>	A ₂₄ G ₂₆ C ₂₁ T ₁₈

<i>Enterococcus faecalis</i>	A ₂₂ G ₂₇ C ₂₀ T ₁₉	<i>Moraxella catarrhalis</i>	A ₂₄ G ₂₆ C ₂₃ T ₁₆
<i>Bacillus megaterium</i>	A ₂₂ G ₂₈ C ₂₀ T ₁₈	<i>Haemophilus influenzae Rd</i>	A ₂₄ G ₂₈ C ₂₀ T ₁₇
<i>Bacillus subtilis</i>	A ₂₂ G ₂₈ C ₂₁ T ₁₇	<i>Chlamydia trachomatis</i>	A ₂₄ G ₂₈ C ₂₁ T ₁₆
<i>Pseudomonas aeruginosa</i>	A ₂₂ G ₂₉ C ₂₃ T ₁₅	<i>Chlamydophila pneumoniae</i>	A ₂₄ G ₂₈ C ₂₁ T ₁₆
<i>Legionella pneumophila</i>	A ₂₂ G ₃₂ C ₂₀ T ₁₆	<i>C. pneumonia AR39</i>	A ₂₄ G ₂₈ C ₂₁ T ₁₆
<i>Mycoplasma pneumoniae</i>	A ₂₃ G ₂₀ C ₁₄ T ₁₆	<i>Pseudomonas putida</i>	A ₂₄ G ₂₉ C ₂₁ T ₁₆
<i>Clostridium botulinum</i>	A ₂₃ G ₂₆ C ₂₀ T ₁₉	<i>Proteus vulgaris</i>	A ₂₄ G ₃₀ C ₂₁ T ₁₅
<i>Enterococcus faecium</i>	A ₂₃ G ₂₆ C ₂₁ T ₁₈	<i>Yersinia pestis</i>	A ₂₄ G ₃₀ C ₂₁ T ₁₅
<i>Acinetobacter calcoaceti</i>	A ₂₃ G ₂₆ C ₂₁ T ₁₉	<i>Yersinia pseudotuberculosis</i>	A ₂₄ G ₃₀ C ₂₁ T ₁₅
<i>Leptospira borgpeterseni</i>	A ₂₃ G ₂₆ C ₂₄ T ₁₅	<i>Clostridium botulinum</i>	A ₂₅ G ₂₄ C ₁₈ T ₂₁
<i>Leptospira interrogans</i>	A ₂₃ G ₂₆ C ₂₄ T ₁₅	<i>Clostridium tetani</i>	A ₂₅ G ₂₅ C ₁₈ T ₂₀
<i>Clostridium perfringens</i>	A ₂₃ G ₂₇ C ₁₉ T ₁₉	<i>Francisella tularensis</i>	A ₂₅ G ₂₅ C ₁₉ T ₁₉
<i>Bacillus anthracis</i>	A ₂₃ G ₂₇ C ₂₀ T ₁₈	<i>Acinetobacter calcoacetic</i>	A ₂₅ G ₂₆ C ₂₀ T ₁₉
<i>Bacillus cereus</i>	A ₂₃ G ₂₇ C ₂₀ T ₁₈	<i>Bacteriodes fragilis</i>	A ₂₅ G ₂₇ C ₁₆ T ₂₂
<i>Bacillus thuringiensis</i>	A ₂₃ G ₂₇ C ₂₀ T ₁₈	<i>Chlamydophila psittaci</i>	A ₂₅ G ₂₇ C ₂₁ T ₁₆
<i>Aeromonas hydrophila</i>	A ₂₃ G ₂₉ C ₂₁ T ₁₆	<i>Borrelia burgdorferi</i>	A ₂₅ G ₂₉ C ₁₇ T ₁₉
<i>Escherichia coli</i>	A ₂₃ G ₂₉ C ₂₁ T ₁₆	<i>Streptobacillus monilifor</i>	A ₂₆ G ₂₆ C ₂₀ T ₁₆
<i>Pseudomonas putida</i>	A ₂₃ G ₂₉ C ₂₁ T ₁₇	<i>Rickettsia prowazekii</i>	A ₂₆ G ₂₈ C ₁₈ T ₁₈
<i>Escherichia coli</i>	A ₂₃ G ₂₉ C ₂₂ T ₁₅	<i>Rickettsia rickettsii</i>	A ₂₆ G ₂₈ C ₂₀ T ₁₆
<i>Shigella dysenteriae</i>	A ₂₃ G ₂₉ C ₂₂ T ₁₅	<i>Mycoplasma mycoides</i>	A ₂₈ G ₂₃ C ₁₆ T ₂₀

The same organism having different base compositions are different strains. Groups of organisms which are highlighted or in italics have the same base compositions in the amplified region. Some of these organisms can be distinguished using multiple primers. For example, *Bacillus anthracis* can be distinguished from *Bacillus cereus* and *Bacillus thuringiensis* using the primer 16S_971-1062 (Table 7). Other primer pairs which produce unique base composition signatures are shown in Table 6 (bold). Clusters containing very similar threat and ubiquitous non-threat organisms (e.g. *anthracis* cluster) are distinguished at high resolution with focused sets of primer pairs. The known biowarfare agents in Table 6 are *Bacillus anthracis*, *Yersinia pestis*, *Francisella tularensis* and *Rickettsia prowazekii*.

Table 7

Organism	16S_971-1062	16S_1228-1310	16S_1100-1188
<i>Aeromonas hydrophila</i>	A ₂₁ G ₂₉ C ₂₂ T ₂₀	A ₂₂ G ₂₇ C ₂₁ T ₁₃	A ₂₃ G ₃₁ C ₂₁ T ₁₅
<i>Aeromonas salmonicida</i>	A ₂₁ G ₂₉ C ₂₂ T ₂₀	A ₂₂ G ₂₇ C ₂₁ T ₁₃	A ₂₃ G ₃₁ C ₂₁ T ₁₅
<i>Bacillus anthracis</i>	A ₂₁ G ₂₇ C ₂₂ T ₂₂	A ₂₄ G ₂₂ C ₁₉ T ₁₈	A ₂₃ G ₂₇ C ₂₀ T ₁₈
<i>Bacillus cereus</i>	A ₂₂ G ₂₇ C ₂₁ T ₂₂	A ₂₄ G ₂₂ C ₁₉ T ₁₈	A ₂₃ G ₂₇ C ₂₀ T ₁₈
<i>Bacillus thuringiensis</i>	A ₂₂ G ₂₇ C ₂₁ T ₂₂	A ₂₄ G ₂₂ C ₁₉ T ₁₈	A ₂₃ G ₂₇ C ₂₀ T ₁₈
<i>Chlamydia trachomatis</i>	A ₂₂ G ₂₆ C ₂₀ T ₂₃	A ₂₄ G ₂₃ C ₁₉ T ₁₆	A ₂₄ G ₂₈ C ₂₁ T ₁₆
<i>Chlamydia pneumoniae AR39</i>	A ₂₆ G ₂₃ C ₂₀ T ₂₂	A ₂₆ G ₂₂ C ₁₆ T ₁₈	A ₂₄ G ₂₈ C ₂₁ T ₁₆
<i>Leptospira borgpetersenii</i>	A ₂₂ G ₂₆ C ₂₀ T ₂₁	A ₂₂ G ₂₅ C ₂₁ T ₁₅	A ₂₃ G ₂₆ C ₂₄ T ₁₅
<i>Leptospira interrogans</i>	A ₂₂ G ₂₆ C ₂₀ T ₂₁	A ₂₂ G ₂₅ C ₂₁ T ₁₅	A ₂₃ G ₂₆ C ₂₄ T ₁₅
<i>Mycoplasma genitalium</i>	A ₂₈ G ₂₃ C ₁₅ T ₂₂	A ₃₀ G ₁₈ C ₁₅ T ₁₉	A ₂₄ G ₁₉ C ₁₂ T ₁₈
<i>Mycoplasma pneumoniae</i>	A ₂₈ G ₂₃ C ₁₅ T ₂₂	A ₂₇ G ₁₉ C ₁₆ T ₂₀	A ₂₃ G ₂₀ C ₁₄ T ₁₆
<i>Escherichia coli</i>	A ₂₂ G ₂₈ C ₂₀ T ₂₂	A ₂₄ G ₂₅ C ₂₁ T ₁₃	A ₂₃ G ₂₉ C ₂₂ T ₁₅
<i>Shigella dysenteriae</i>	A ₂₂ G ₂₈ C ₂₁ T ₂₁	A ₂₄ G ₂₅ C ₂₁ T ₁₃	A ₂₃ G ₂₉ C ₂₂ T ₁₅
<i>Proteus vulgaris</i>	A ₂₃ G ₂₆ C ₂₂ T ₂₁	A ₂₆ G ₂₄ C ₁₉ T ₁₄	A ₂₄ G ₃₀ C ₂₁ T ₁₅
<i>Yersinia pestis</i>	A ₂₄ G ₂₅ C ₂₁ T ₂₂	A ₂₅ G ₂₄ C ₂₀ T ₁₄	A ₂₄ G ₃₀ C ₂₁ T ₁₅
<i>Yersinia pseudotuberculosis</i>	A ₂₄ G ₂₅ C ₂₁ T ₂₂	A ₂₅ G ₂₄ C ₂₀ T ₁₄	A ₂₄ G ₃₀ C ₂₁ T ₁₅
<i>Francisella tularensis</i>	A ₂₀ G ₂₅ C ₂₁ T ₂₃	A ₂₃ G ₂₆ C ₁₇ T ₁₇	A ₂₄ G ₂₆ C ₁₉ T ₁₉
<i>Rickettsia prowazekii</i>	A ₂₁ G ₂₆ C ₂₄ T ₂₅	A ₂₄ G ₂₃ C ₁₆ T ₁₉	A ₂₆ G ₂₈ C ₁₈ T ₁₈
<i>Rickettsia rickettsii</i>	A ₂₁ G ₂₆ C ₂₅ T ₂₄	A ₂₄ G ₂₄ C ₁₇ T ₁₇	A ₂₆ G ₂₈ C ₂₀ T ₁₆

The sequence of *B. anthracis* and *B. cereus* in region 16S_971 is shown below. Shown in bold is the single base difference between the two species that can be detected using the 5 methods of the present invention. *B. anthracis* has an ambiguous base at position 20.

*B.anthraxis*_16S_971

GCGAAGAACCUUACCAGGUNUUGACAUCCUCUGACAACCCUAGAGAUAGGGCUUC
UCCUUCGGGAGCAGAGUGACAGGUGGUGCAUGGUU (SEQ ID NO:1)

*B.cereus*_16S_971

10 GCGAAGAACCUUACCAGGUCUUGACAUCCUCUGAAAACCCUAGAGAUAGGGCUUC
UCCUUCGGGAGCAGAGUGACAGGUGGUGCAUGGUU (SEQ ID NO:2)

Example 6: ESI-TOF MS of sspE 56-mer Plus Calibrant

The mass measurement accuracy that can be obtained using an internal mass standard in the ESI-MS study of PCR products is shown in Fig.8. The mass standard was a 20-mer phosphorothioate oligonucleotide added to a solution containing a 56-mer PCR product from the *B. anthracis* spore coat protein sspE. The mass of the expected PCR product distinguishes *B. anthracis* from other species of Bacillus such as *B. thuringiensis* and *B. cereus*.

Example 7: *B. anthracis* ESI-TOF Synthetic 16S_1228 Duplex

An ESI-TOF MS spectrum was obtained from an aqueous solution containing 5 μ M each of synthetic analogs of the expected forward and reverse PCR products from the nucleotide 1228 region of the *B. anthracis* 16S rRNA gene. The results (Fig. 9) show that the molecular weights of the forward and reverse strands can be accurately determined and easily distinguish the two strands. The $[M-21H]^+{}^{21-}$ and $[M-20H]^+{}^{20-}$ charge states are shown.

Example 8: ESI-FTICR-MS of Synthetic *B. anthracis* 16S_1337 46 Base Pair Duplex

An ESI-FTICR-MS spectrum was obtained from an aqueous solution containing 5 μ M each of synthetic analogs of the expected forward and reverse PCR products from the nucleotide 1337 region of the *B. anthracis* 16S rRNA gene. The results (Fig. 10) show that the molecular weights of the strands can be distinguished by this method. The $[M-16H]^+{}^{16-}$ through $[M-10H]^+{}^{10-}$ charge states are shown. The insert highlights the resolution that can be realized on the FTICR-MS instrument, which allows the charge state of the ion to be determined from the mass difference between peaks differing by a single ^{13}C substitution.

Example 9: ESI-TOF MS of 56-mer Oligonucleotide from saspB Gene of *B. anthracis* with Internal Mass Standard

ESI-TOF MS spectra were obtained on a synthetic 56-mer oligonucleotide (5 μ M) from the saspB gene of *B. anthracis* containing an internal mass standard at an ESI of 1.7 μ L/min as a function of sample consumption. The results (Fig. 11) show that the signal to noise is improved as more scans are summed, and that the standard and the product are visible after only 100 scans.

Example 10: ESI-TOF MS of an Internal Standard with Tributylammonium (TBA)-trifluoroacetate (TFA) Buffer

An ESI-TOF-MS spectrum of a 20-mer phosphorothioate mass standard was obtained following addition of 5 mM TBA-TFA buffer to the solution. This buffer strips charge from the oligonucleotide and shifts the most abundant charge state from $[M-8H]^8-$ to $[M-3H]^3-$ (Fig. 12).

5 Example 11: Master Database Comparison

The molecular masses obtained through Examples 1-10 are compared to molecular masses of known bioagents stored in a master database to obtain a high probability matching molecular mass.

10 Example 12: Master Data Base Interrogation over the Internet

The same procedure as in Example 11 is followed except that the local computer did not store the Master database. The Master database is interrogated over an internet connection, searching for a molecular mass match.

15 Example 13: Master Database Updating

The same procedure as in example 11 is followed except the local computer is connected to the internet and has the ability to store a master database locally. The local computer system periodically, or at the user's discretion, interrogates the Master database, synchronizing the local master database with the global Master database. This provides the
20 current molecular mass information to both the local database as well as to the global Master database. This further provides more of a globalized knowledge base.

Example 14: Global Database Updating

The same procedure as in example 13 is followed except there are numerous such local
25 stations throughout the world. The synchronization of each database adds to the diversity of information and diversity of the molecular masses of known bioagents.

Example 15: Demonstration of Detection and Identification of Five Species of Bacteria in a Mixture

30 Broad range intelligent primers were chosen following analysis of a large collection of curated bacterial 16S rRNA sequences representing greater than 4000 species of bacteria. Examples of primers capable of priming from greater than 90% of the organisms in the collection include, but are not limited to, those exhibited in Table 8 wherein Tp = 5'propynylated uridine and Cp = 5'propynylated cytidine.

Table 8
Intelligent Primer Pairs for Identification of Bacteria

Primer Pair Name	Forward Primer Sequence	Forward SEQ ID NO:	Reverse Primer Sequence	Reverse SEQ ID NO:
16S_EC_107 7_1195	GTGAGATGTTGGGTAAAGTCCC GTAACGAG	8	GACGTCATCCCCACCTTCCTC	9
16S_EC_108 2_1197	ATGTTGGGTAAAGTCCGCAAC GAG	10	TGACGTCATCCCCACCTTCCT C	11
16S_EC_109 0_1196	TTAAGTCCCGCAACGATCGCAA	12	TGACGTCATCCCCACCTTCCTC	13
16S_EC_122 2_1323	GCTACACACGTGCTACAATG	14	CGAGTTGCAGACTGCGATCCG	15
16S_EC_133 2_1407	AAGTCGGAATCGCTAGTAATCG	16	GACGGGCGGTGTGTACAAG	17
16S_EC_30_ 126	TGAACGCTGGTGGCATGCTTAA CAC	18	TACGCATTACTCACCCGTCCGC	19
16S_EC_38_ 120	GTGGCATGCCTAATACATGCAA GTCG	20	TTACTCACCCGTCCGCCGCT	21
16S_EC_49_ 120	TAACACATGCAAGTCGAACG	22	TTACTCACCCGTCCGCC	23
16S_EC_683 795	GTGTAGCGGTGAAATGCG	24	GTATCTAATCCTGTTGCTCCC	25
16S_EC_713 809	AGAACACCGATGGCGAAGGC	26	CGTGGACTACCAGGGTATCTA	27
16S_EC_785 897	GGATTAGAGACCCTGGTAGTCC	28	GGCCGTACTCCCCAGGCG	29
16S_EC_785 _897_2	GGATTAGATACCCTGGTAGTCC ACGC	30	GGCCGTACTCCCCAGGCG	31
16S_EC_789 894	TAGATACCCTGGTAGTCCACGC	32	CGTACTCCCCAGGCG	33
16S_EC_960 1073	TTGATGCAACGCGAAGAACCT	34	ACGAGCTGACGACAGCCATG	35
16S_EC_969 1078	ACGCGAAGAACCTTACC	36	ACGACACGAGCTGACGAC	37
23S_EC_182 6_1924	CTGACACCTGCCCCGGTGC	38	GACCGTTATAGTTACGGCC	39
23S_EC_264 5_2761	TCTGTCCCTAGTACGAGAGGAC CGG	40	TGCTTAGATGCTTTCAGC	41
23S_EC_264 5_2767	CTGTCCCTAGTACGAGAGGACC GG	42	GTTTCATGCTTAGATGCTTCA GC	43
23S_EC_493 _571	GGGGAGTGAAAGAGATCCTGAA ACCG	44	ACAAAAGGTACGCCGTCACCC	45
23S_EC_493 _571_2	GGGGAGTGAAAGAGATCCTGAA ACCG	46	ACAAAAGGCACGCCATCACCC	47
23S_EC_971 1077	CGAGAGGGAAACAACCCAGACC	48	TGGCTGCTTCTAAGCCAAC	49
INFB_EC_13 65_1467	TGCTCGTGGTGCACAAGTAACG GATATTA	50	TGCTGCTTTCGCATGGTTAATT GCTTCAA	51
RPOC_EC_10 18_1124	CAAACTTATTAGGTAAGCGTG TTGACT	52	TCAAGCGCCATTTCTTTGGTA AACCACAT	53
RPOC_EC_10 18_1124_2	CAAACTTATTAGGTAAGCGTG TTGACT	54	TCAAGCGCCATCTCTTTCGGTA ATCCACAT	55
RPOC_EC_11 4_232	TAAGAAGCCGGAACCATCAAC TACCG	56	GGCGCTGTACTTACCGCAC	57

RPOC_EC_21 78_2246	TGATTCTGGTGCCCGTGGT	58	TTGGCCATCAGGCCACGCATAC	59
RPOC_EC_21 78_2246_2	TGATTCCGGTGCCCGTGGT	60	TTGGCCATCAGACCACGCATAC	61
RPOC_EC_22 18_2337	CTGGCAGGTATGCGTGGTCTGA TG	62	CGCACCGTGGGTTGAGATGAAG TAC	63
RPOC_EC_22 18_2337_2	CTTGCTGGTATGCGTGGTCTGA TG	64	CGCACCATGCGTAGAGATGAAG TAC	65
RPOC_EC_80 8_889	CGTCGGGTGATTAACCGTAACA ACCG	66	GTTTTTCGTTGCGTACGATGAT GTC	67
RPOC_EC_80 8_891	CGTCGTGTAATTAACCGTAACA ACCG	68	ACGTTTTTCGTTTGAACGATA ATGCT	69
RPOC_EC_99 3_1059	CAAAGGTAAGCAAGGTCGTTTC CGTCA	70	CGAACGGCCTGAGTAGTCAACA CG	71
RPOC_EC_99 3_1059_2	CAAAGGTAAGCAAGGACGTTTC CGTCA	72	CGAACGGCCAGAGTAGTCAACA CG	73
TUFB_EC_23 9_303	TAGACTGCCCAGGACACGCTG	74	GCCGTCCATCTGAGCAGCACC	75
TUFB_EC_23 9_303_2	TTGACTGCCCAGGTCACGCTG	76	GCCGTCCATTTGAGCAGCACC	77
TUFB_EC_97 6_1068	AACTACCGTCCGCAGTTCTACT TCC	78	GTTGTCGCCAGGCATAACCATT TC	79
TUFB_EC_97 6_1068_2	AACTACCGTCCTCAGTTCTACT TCC	80	GTTGTCACCAGGCATTACCATT TC	81
TUFB_EC_98 5_1062	CCACAGTTCTACTTCCGTACTA CTGACG	82	TCCAGGCATTACCATTCTACT CCTTCTGG	83
RPLB_EC_65 0_762	GACCTACAGTAAGAGGTTCTGT AATGAACC	84	TCCAAGTGCTGGTTTACCCCAT GG	85
RPLB_EC_68 8_757	CATCCACACGGTGGTGGTGAAG G	86	GTGCTGGTTTACCCCATGGAGT	87
RPOC_EC_10 36_1126	CGTGTGACTATTGCGGGCGTT CAG	88	ATTCAAGAGCCATTCTTTTGG TAAACCAC	89
RPOB_EC_37 62_3865	TCAACAACCTCTTGAGGTAAA GCTCAGT	90	TTTCTTGAAGATATGAGCTGC TCCGTAAG	91
RPLB_EC_68 8_771	CATCCACACGGTGGTGGTGAAG G	92	TGTTTTGTATCCAAGTGCTGGT TTACCCC	93
VALS_EC_11 05_1218	CGTGGCGGCGTGTTATCGA	94	CGGTACGAACTGGATGTCGCCG TT	95
RPOB_EC_18 45_1929	TATCGCTCAGGCGAACTCCAAC	96	GCTGGATTGCGCTTTGCTACG	97
RPLB_EC_66 9_761	TGTAATGAACCCTAATGACCAT CCACACGG	98	CCAAGTGCTGGTTTACCCCATG GAGTA	99
RPLB_EC_67 1_762	TAATGAACCCTAATGACCATCC ACACGGTG	100	TCCAAGTGCTGGTTTACCCCAT GGAG	101
RPOB_EC_37 75_3858	CTTGGAGGTAAGTCTCATTTTG GTGGGCA	102	CGTATAAGCTGCACCATAAGCT TGTAATGC	103
VALS_EC_18 33_1943	CGACGCGCTGCGCTTCAC	104	GCGTTCCACAGCTTGTTGCAGA AG	105
RPOB_EC_13 36_1455	GACCACCTCGGCAACCGT	106	TTCGCTCTCGGCCTGGCC	107
TUFB_EC_22 5_309	GCACTATGCACACGTAGATTGT CCTGG	108	TATAGCACCATCCATCTGAGCG GCAC	109
DNAK_EC_42 8_522	CGGCGTACTTCAACGACAGCCA	110	CGCGGTGCGCTCGTTGATGA	111

VALS_EC_19 20_1970	CTTCTGCAACAAGCTGTGGAAC GC	112	TCGCAGTTCATCAGCACGAAGC G	113
TUFB_EC_75 7_867	AAGACGACCTGCACGGGC	114	GCGCTCCACGTCTTCACGC	115
23S_EC_264 6_2765	CTGTTCTTAGTACGAGAGGACC	116	TTCGTGCTTAGATGCTTTTCAG	117
16S_EC_969 _1078_3P	ACGCGAAGAACCTTACpC	118	ACGACACGAGCpTpGACGAC	119
16S_EC_972 _1075_4P	CGAAGAACpCpTTACC	120	ACACGAGCpTpGAC	121
16S_EC_972 _1075	CGAAGAACCTTACC	122	ACACGAGCTGAC	123
23S_EC_- 347_59	CCTGATAAGGGTGAGGTCG	124	ACGTCCTTCATCGCCTCTGA	125
23S_EC_- 7_450	GTTGTGAGGTTAAGCGACTAAG	126	CTATCGGTCAAGTCAGGAGTAT	127
23S_EC_- 7_910	GTTGTGAGGTTAAGCGACTAAG	128	TTGCATCGGGTTGGTAAGTC	129
23S_EC_430 _1442	ATACTCCTGACTGACCGATAG	130	AACATAGCCTTCTCCGTCC	131
23S_EC_891 _1931	GACTTACCAACCCGATGCAA	132	TACCTTAGGACCGTTATAGTTA CG	133
23S_EC_142 4_2494	GGACGGAGAAGGCTATGTT	134	CCAAACACCGCCGTCGATAT	135
23S_EC_190 8_2852	CGTAACTATAACGGTCCTAAGG TA	136	GCTTACACACCCGGCCTATC	137
23S_EC_247 5_3209	ATATCGACGGCGGTGTTTGG	138	GCGTGACAGGCAGGTATTC	139
16S_EC_- 60_525	AGTCTCAAGAGTGAACACGTAA	140	GCTGCTGGCACGGAGTTA	141
16S_EC_326 _1058	GACACGGTCCAGACTCCTAC	142	CCATGCAGCACCTGTCTC	143
16S_EC_705 _1512	GATCTGGAGGAATACCGGTG	144	ACGGTTACCTTGTTACGACT	145
16S_EC_126 8_1775	GAGAGCAAGCGGACCTCATA	146	CCTCCTGCGTGCAAAGC	147
GROL_EC_94 1_1060	TGGAAGATCTGGGTCAAGC	148	CAATCTGCTGACGGATCTGAGC	149
INFB_EC_11 03_1191	GTCGTGAAAACGAGCTGGAAGA	150	CATGATGGTCACAACCGG	151
HFLB_EC_10 82_1168	TGGCGAACCTGGTGAACGAAGC	152	CTTTCGCTTTCTCGAACTCAAC CAT	153
INFB_EC_19 69_2058	CGTCAGGGTAAATCCGTGAAG TTAA	154	AACTTCGCCTTCGGTCATGTT	155
GROL_EC_21 9_350	GGTGAAAGAAGTTGCCTCTAAA GC	156	TTCAGGTCCATCGGGTTCATGC C	157
VALS_EC_11 05_1214	CGTGGCGGCGTGTTATCGA	158	ACGAACTGGATGTCGCCGTT	159
16S_EC_556 _700	CGGAATTACTGGGCGTAAAG	160	CGCATTTACCGCTACAC	161
RPOC_EC_12 56_1315	ACCCAGTGCTGCTGAACCGTGC	162	GTTCAAATGCCTGGATACCCA	163
16S_EC_774 _894	GGGAGCAAACAGGATTAGATAC	164	CGTACTCCCCAGGCG	165
RPOC_EC_15 84_1643	TGGCCCGAAAGAAGCTGAGCG	166	ACGCGGGCATGCAGAGATGCC	167
16S_EC_108 2_1196	ATGTTGGGTTAAGTCCCGC	168	TGACGTCATCCCCACCTTCC	169

16S_EC_138 9_1541	CTGTGACACACCGCCGTC	170	AAGGAGGTGATCCAGCC	171
16S_EC_130 3_1407	CGGATTGGAGTCTGCAACTCG	172	GACGGGCGGTGTGTACAAG	173
23S_EC_23_ 130	GGTGGATGCCTTGGC	174	GGGTTTCCCCATTCCG	175
23S_EC_187 256	GGGAACTGAAACATCTAAGTA	176	TTCGCTCGCCGCTAC	177
23S_EC_160 2_1703	TACCCCAAACCGACACAGG	178	CCTTCTCCCGAAGTTACG	179
23S_EC_168 5_1842	CCGTAACCTTCGGGAGAAGG	180	CACCGGGCAGGCGTC	181
23S_EC_182 7_1949	GACGCCTGCCCGGTGC	182	CCGACAAGGAATTTCGCTACC	183
23S_EC_243 4_2511	AAGGTACTCCGGGGATAACAGG C	184	AGCCGACATCGAGGTGCCAAAC	185
23S_EC_259 9_2669	GACAGTTCGGTCCCTATC	186	CCGGTCTCTCGTACTA	187
23S_EC_265 3_2758	TAGTACGAGAGGACCGG	188	TTAGATGCTTTCAGCACTTATC	189
23S_BS_- 68_21	AAACTAGATAACAGTAGACATC AC	190	GTGCGCCCTTTCTAACTT	191
16S_EC_8_3 58	AGAGTTTGATCATGGCTCAG	192	ACTGCTGCCTCCCGTAG	193
16S_EC_314 575	CACTGGAAGTGAACACGG	194	CTTACGCCCAGTAATTCCG	195
16S_EC_518 795	CCAGCAGCCGCGGTAATAC	196	GTATCTAATCCTGTTGCTCCC	197
16S_EC_683 985	GTGTAGCGGTGAAATGCG	198	GGTAAGGTTCTTCGCGTTG	199
16S_EC_937 1240	AAGCGGTGGAGCATGTGG	200	ATTGTAGCACGTGTGTAGCCC	201
16S_EC_119 5_1541	CAAGTCATCATGGCCCTTA	202	AAGGAGGTGATCCAGCC	203
16S_EC_8_1 541	AGAGTTTGATCATGGCTCAG	204	AAGGAGGTGATCCAGCC	205
23S_EC_183 1_1936	ACCTGCCAGTGCTGGAAG	206	TCGCTACCTTAGGACCGT	207
16S_EC_138 7_1513	GCCTTGACACACCTCCCGTC	208	CACGGCTACCTTGTTACGAC	209
16S_EC_139 0_1505	TTGTACACACCGCCCGTCATAC	210	CCTTGTTACGACTTCACCCC	211
16S_EC_136 7_1506	TACGGTGAATACGTTCCCGGG	212	ACCTTGTTACGACTTCACCCCA	213
16S_EC_804 929	ACCACGCCGTAAACGATGA	214	CCCCCGTCAATTCCTTTGAGT	215
16S_EC_791 904	GATACCCTGGTAGTCCACACCG	216	GCCTTGCGACCGTACTCCC	217
16S_EC_789 899	TAGATACCCTGGTAGTCCACGC	218	GCGACCGTACTCCCCAGG	219
16S_EC_109 2_1195	TAGTCCCGCAACGAGCGC	220	GACGTCATCCCCACCTTCCTCC	221
23S_EC_258 6_2677	TAGAACGTGCGAGACAGTTG	222	AGTCCATCCCGGTCCTCTCG	223
HEXAMER_EC 61_362	GAGGAAAGTCCGGGCTC	224	ATAAGCCGGGTTCTGTGCG	225
RNASEP_BS_ 43_384	GAGGAAAGTCCATGCTCGC	226	GTAAGCCATGTTTTGTTCCATC	227
RNASEP_EC_ 43_384	GAGGAAAGTCCGGGCTC	228	ATAAGCCGGGTTCTGTGCG	229

61_362				
YAED_TRNA_31_379 ALA- RRNH_EC_51 3_49	GCGGGATCCTCTAGAGGTGTTA AATAGCCTGGCAG	230	GCGGGATCCTCTAGAAGACCTC CTGCGTGCAAAGC	231
RNASEP_SA_31_379	GAGGAAAGTCCATGCTCAC	232	ATAAGCCATGTTCTGTTCATC	233
16S_EC_108 2_1541	ATGTTGGGTAAAGTCCCGC	234	AAGGAGGTGATCCAGCC	235
16S_EC_556 795	CGGAATTACTGGGCGTAAAG	236	GTATCTAATCCTGTTGCTCCC	237
16S_EC_108 2_1196_10G	ATGTTGGGTAAAGTCCCGC	238	TGACGTCATGCCACCTTCC	239
16S_EC_108 2_1196_10G 11G	ATGTTGGGTAAAGTCCCGC	240	TGACGTCATGGCCACCTTCC	241
TRNA_ILERR NH_ASPRRNH EC_32_41	GCGGGATCCTCTAGACCTGATA AGGGTGAGGTCG	242	GCGGGATCCTCTAGAGCGTGAC AGGCAGGTATTC	243
16S_EC_969 1407	ACGCGAAGAACCTTACC	244	GACGGGCGGTGTGTACAAG	245
16S_EC_683 1323	GTGTAGCGGTGAAATGCG	246	CGAGTTGCAGACTGCGATCCG	247
16S_EC_49 894	TAACACATGCAAGTCGAACG	248	CGTACTCCCCAGGCG	249
16S_EC_49 1078	TAACACATGCAAGTCGAACG	250	ACGACACGAGCTGACGAC	251
CYA_BA_134 9_1447	ACAACGAAGTACAATACAAGAC	252	CTTCTACATTTTTAGCCATCAC	253
16S_EC_109 0_1196_2	TTAAGTCCCGCAACGAGCGCAA	254	TGACGTCATCCCCACCTTCCTC	255
16S_EC_405 527	TGAGTGATGAAGGCCTTAGGGT TGTAAG	256	CGGCTGCTGGCAGGAAGTTAG	257
GROL_EC_49 6_596	ATGGACAAGGTTGGCAAGGAAG G	258	TAGCCGCGGTGCAATTGCAT	259
GROL_EC_51 1_593	AAGGAAGGCGTGATCACC GTTGAAGA	260	CCGCGGTGCAATTGCATGCCTT C	261
VALS_EC_18 35_1928	ACGCGCTGCGCTTAC	262	TTGCAGAAGTTGCGGTAGCC	263
RPOB_EC_13 34_1478	TCGACCACCTGGGCAACC	264	ATCAGGTCGTGCGGCATCA	265
DNAK_EC_42 0_521	CACGGTGCCGGCGTACT	266	GCGGTCGGCTCGTTGATGAT	267
RPOB_EC_37 76_3853	TTGGAGGTAAGTCTCATTTTGG TGG	268	AAGCTGCACCATAAGCTTGTAATGC	269
RPOB_EC_38 02_3885	CAGCGTTTCGGCGAAATGGA	270	CGACTTGACGGTTAACATTTCC TG	271
RPOB_EC_37 99_3888	GGGCAGCGTTTCGGCGAAATGGA A	272	GTCCGACTTGACGGTCAACATT TCCTG	273
RPOC_EC_21 46_2245	CAGGAGTCGTTCAACTCGATCT ACATGAT	274	ACGCCATCAGGCCACGCAT	275
ASPS_EC_40 5_538	GCACAACCTGCGGCTGCG	276	ACGGCAGGAGGTAGTCGC	277
RPOC_EC_13 74_1455	CGCCGACTTCGACGGTGACC	278	GAGCATCAGCGTGCGTGCT	279
TUFB_EC_95 7_1058	CCACAGCCGTTCTTCAACAAC T	280	GGCATCACCATTTCCTTGTCCT TCG	281

16S_EC_7_1 22	GAGAGTTTGATCCTGGCTCAGA ACGAA	282	TGTTACTCACCCTCTGCCACT	283
VALS_EC_61 0 727	ACCGAGCAAGGAGACCAGC	284	TATAACGCACATCGTCAGGGTG A	285

For evaluation in the laboratory, five species of bacteria were selected including three γ -proteobacteria (*E. coli*, *K. pneumoniae* and *P. aeruginosa*) and two low G+C gram positive bacteria (*B. subtilis* and *S. aureus*). The identities of the organisms were not revealed to the laboratory technicians.

Bacteria were grown in culture, DNA was isolated and processed, and PCR performed using standard protocols. Following PCR, all samples were desalted, concentrated, and analyzed by Fourier Transform Ion Cyclotron Resonance (FTICR) mass spectrometry. Due to the extremely high precision of the FTICR, masses could be measured to within 1 Da and unambiguously deconvoluted to a single base composition. The measured base compositions were compared with the known base composition signatures in our database. As expected when using broad range survey 16S primers, several phylogenetic near-neighbor organisms were difficult to distinguish from our test organisms. Additional non-ribosomal primers were used to triangulate and further resolve these clusters.

An example of the use of primers directed to regions of RNA polymerase B (rpoB) is shown in Figure 19. This gene has the potential to provide broad priming and resolving capabilities. A pair of primers directed against a conserved region of rpoB provided distinct base composition signatures that helped resolve the tight enterobacteriae cluster. Joint probability estimates of the signatures from each of the primers resulted in the identification of a single organism that matched the identity of the test sample. Therefore a combination of a small number of primers that amplify selected regions of the 16S ribosomal RNA gene and a few additional primers that amplify selected regions of protein encoding genes provide sufficient information to detect and identify all bacterial pathogens.

Example 16: Detection of *Staphylococcus aureus* in Blood Samples

Blood samples in an analysis plate were spiked with genomic DNA equivalent of 10^3 organisms/ml of *Staphylococcus aureus*. A single set of 16S rRNA primers was used for amplification. Following PCR, all samples were desalted, concentrated, and analyzed by Fourier Transform Ion Cyclotron Resonance (FTICR) mass spectrometry. In each of the spiked wells, strong signals were detected which are consistent with the expected BCS of the *S. aureus* amplicon (Figure 20). Furthermore, there was no robotic carryover or contamination in any of

the blood only or water blank wells. Methods similar to this one will be applied for other clinically relevant samples including, but not limited to: urine and throat or nasal swabs.

Example 17: Detection and Serotyping of Viruses

5 The virus detection capability of the present invention was demonstrated in collaboration with Naval health officers using adenoviruses as an example.

All available genomic sequences for human adenoviruses available in public databases were surveyed. The hexon gene was identified as a candidate likely to have broad specificity across all serotypes. Four primer pairs were selected from a group of primers designed to yield
10 broad coverage across the majority of the adenoviral strain types (Table 9) wherein Tp = 5'propynylated uridine and Cp = 5'propynylated cytidine.

Table 9

Intelligent Primer Pairs for Serotyping of Adenoviruses

Primer Pair Name	Forward Primer Sequence	Forward SEQ ID NO:	Reverse Primer Sequence	Reverse SEQ ID NO:
HEX_HAD7+4+2 1 934 995	AGACCCAATTACATTGGCTT	286	CCAGTGCTGTGTAGTACAT	287
HEX_HAD7+4+2 1 976 1050	ATGTACTACAACAGTACTGG	288	CAAGTCAACCACAGCATTCA	289
HEX_HAD7+4+2 1 970 1059	GGGCTTATGTACTACAACAG	290	TCTGTCTTGCAAGTCAACCAC	291
HEX_HAD7+3_7 71 827	GGAATTTTTTGATGGTAGAGA	292	TAAAGCACAAATTCAGGCG	293
HEX_HAD4+16_ 746 848	TAGATCTGGCTTTCTTTGAC	294	ATATGAGTATCTGGAGTCTGC	295
HEX_HAD7_509 578	GGAAAGACATTACTGCAGACA	296	CCAACTTGAGGCTCTGGCTG	297
HEX_HAD4_121 6 1289	ACAGACACTTACCAGGGTG	298	ACTGTGGTGTCTCTTTGTC	299
HEX_HAD21_51 5 567	TCACTAAAGACAAAGGTCTTC	300	GGCTTCGCCGTCTGTAATTC	301
HEX_HAD_1342 1469	CGGATCCAAGCTAATCTTTGG	302	GGTATGTACTCATAGGTGTTG GTG	303
HEX_HAD7+4+2 1 934 995P	AGACpCpCAATTpACpATpTGG CTT	304	CpCpAGTGCTGTpTpGTAGTA CAT	305
HEX_HAD7+4+2 1 976 1050P	ATpGTpACTpACAACAGTACpT pGG	306	CAAGTpCpAACCACAGCATpT pCA	307
HEX_HAD7+4+2 1 970 1059P	GGGCpTpTATpGTpACTACAAC pAG	308	TCTGTpCpTTGCAAGTpCpAA CCAC	309
HEX_HAD7+3_7 71 827P	GGAATTpTpTpTpTGATGGTAG AGA	310	TAAAGCACAAATpTpTpCpAGG CG	311
HEX_HAD4+16_ 746 848P	TAGATCTGGCTpTpTpCpTTTG AC	312	ATATGAGTATpCpTpGGAGTp CpTGC	313
HEX_HAD_1342 1469P	CGGATpCCAAGCpTAATCpTpT TGG	314	GGTATGTACTCATAGGTGTpT pGGTG	315
HEX_HAD7+21+	AACAGACCCAATTACATTGGCT	316	GAGGCACTTGTATGTGAAAG	317

3_931_1645	T		G	
HEX_HAD4+2_9 25_1469	ATGCCTAACAGACCCAATTACA T	318	TTCATGTAGTCGTAGGTGTTG G	319
HEX_HAD7+21+ 3_384_953	CGCGCCTAATACATCTCAGTGG AT	320	AAGCCAATGTAATTGGGTCTG TT	321
HEX_HAD4+2_3 45_947	CTACTCTGGCACTGCCTACAAC	322	ATGTAATTGGGTCTGTTAGGC AT	323
HEX_HAD2_772 865	CAATCCGTTCTGGTTCCGGATG AA	324	CTTGCCGGTCGTTCAAAGAGG TAG	325
HEX_HAD7+4+2 1_73_179	AGTCCGGGTCTGGTGCAG	326	CGGTCCGGTGGTCACATC	327
HEX_HAD7+4+2 1_1_54	ATGGCCACCCCATCGATG	328	CTGTCCGGCGATGTGCATG	329
HEX_HAD7+4+2 1_1612_1718	GGTCGTTATGTGCCTTTCCACA T	330	TCCTTTCTGAAGTTCCTCA TAGG	331
HEX_HAD7+4+2 1_2276_2368	ACAACATTGGCTACCAGGGCTT	332	CCTGCCTGCTCATAGGCTGGA AGTT	333

These primers also served to clearly distinguish those strains responsible for most disease (types 3, 4, 7 and 21) from all others. DNA isolated from field samples known to contain adenoviruses were tested using the hexon gene PCR primers, which provided unambiguous strain identification for all samples. A single sample was found to contain a mixture of two viral DNAs belonging to strains 7 and 21.

Test results (Figure 21) showed perfect concordance between predicted and observed base composition signatures for each of these samples. Classical serotyping results confirmed each of these observations. Processing of viral samples directly from collection material such as throat swabs rather than from isolated DNA, will result in a significant increase in throughput, eliminating the need for virus culture.

Example 18: Broad Rapid Detection and Strain Typing of Respiratory Pathogens for Epidemic Surveillance

Genome preparation: Genomic materials from culture samples or swabs were prepared using a modified robotic protocol using DNeasy™ 96 Tissue Kit, Qiagen). Cultures of *Streptococcus pyogenes* were pelleted and transferred to a 1.5 mL tube containing 0.45 g of 0.7 mm Zirconia beads (Biospec Products, Inc.). Cells were lysed by shaking for 10 minutes at a speed of 19 1/s using a MM300 Vibration Mill (Retsch, Germany). The samples were centrifuged for 5 min and the supernatants transferred to deep well blocks and processed using the manufacture's protocol and a Qiagen 8000 BioRobot.

PCR: PCR reactions were assembled using a Packard MPII liquid handling platform and were performed in 50 µL volume using 1.8 units each of Platinum Taq (Invitrogen) and Hotstart PFU Turbo (Stratagene) polymerases. Cycling was performed on a DNA Engine Dyad

(MJ Research) with cycling conditions consisting of an initial 2 min at 95°C followed by 45 cycles of 20 s at 95°C, 15 s at 58°C, and 15 s at 72°C.

Broad-range primers: PCR primer design for base composition analysis from precise mass measurements is constrained by an upper limit where ionization and accurate deconvolution can be achieved. Currently, this limit is approximately 140 base pairs. Primers designed to broadly conserved regions of bacterial ribosomal RNAs (16 and 23S) and the gene encoding ribosomal protein L3 (rpoC) are shown in Table 10.

Table 10
Broad Range Primer Pairs

Target Gene	Direction	Primer	SEQ ID NO	Length of Amplicon
16S_1	F	GGATTAGAGACCTGGTAGTCC	334	116
16S_1	R	GGCCGTACTCCCCAGGCG	335	116
16S_2	F	TTCGATGCAACGCGAAGAACCT	336	115
16S_2	R	ACGAGCTGACGACAGCCATG	337	115
23S	F	TCTGTCCCTAGTACGAGAGGACCGG	338	118
23S	R	TGCTTAGATGCTTTCAGC	339	118
rpoC	F	CTGGCAGGTATGCGTGGTCTGATG	340	121
rpoC	R	CGCACC GTGGGTTGAGATGAAGTAC	341	121

10

Emm-typing primers: The allelic profile of a GAS strain by Multilocus Sequencing Technique (MLST) can be obtained by sequencing the internal fragments of seven housekeeping genes. The nucleotide sequences for each of these housekeeping genes, for 212 isolates of GAS (78 distinct emm types), are available (www.mlst.net). This corresponds to one hundred different allelic profiles or unique sequence types, referred to by Enright et al. as ST1-ST100 (Enright et al., *Infection and Immunity*, 2001, 69, 2416-2427). For each sequence type, we created a virtual transcript by concatenating sequences appropriate to their allelic profile from each of the seven genes. MLST primers were designed using these sequences and were constrained to be within each gene loci. Twenty-four primer pairs were initially designed and tested against the sequenced GAS strain 700294. A final subset of six primer pairs Table 11 was chosen based on a theoretical calculation of minimal number of primer pairs that maximized resolution of between emm types.

25

Table 11
Drill-Down Primer Pairs Used in Determining emm-type

Target Gene	Direction	Primer	SEQ ID NO	Length of Amplicon
<i>gki</i>	F	GGGGATTTCAGCCATCAAAGCAGCTATTGA C	342	116
<i>gki</i>	R	CCAACCTTTTCCACAACAGAATCAGC	343	116
<i>gtr</i>	F	CCTTACTTCGAACTATGAATCTTTTGGAA G	344	115
<i>gtr</i>	R	CCCATTTTTTTCACGCATGCTGAAAATATC	345	115
<i>murI</i>	F	CGCAAAAAAATCCAGCTATTAGC	346	118
<i>murI</i>	R	AAACTATTTTTTTAGCTATACTCGAACAC	347	118
<i>mutS</i>	F	ATGATTACAATTCAAGAAGGTCGTCACGC	348	121
<i>mutS</i>	R	TTGGACCTGTAATCAGCTGAATACTGG	349	121
<i>xpt</i>	F	GATGACTTTTTAGCTAATGGTCAGGCAGC	350	122
<i>xpt</i>	R	AATCGACGACCATCTTGGAAGATTCTC	351	122
<i>yqiL</i>	F	GCTTCAGGAATCAATGATGGAGCAG	352	119
<i>yqiL</i>	R	GGGTCTACACCTGCACTTGCATAAC	353	119

Microbiology: GAS isolates were identified from swabs on the basis of colony morphology and beta-hemolysis on blood agar plates, gram stain characteristics, susceptibility to bacitracin, and positive latex agglutination reactivity with group A-specific antiserum.

Sequencing: Bacterial genomic DNA samples of all isolates were extracted from freshly grown GAS strains by using QIAamp DNA Blood Mini Kit (Qiagen, Valencia, CA) according to the procedures described by the manufacture. Group A streptococcal cells were subjected to PCR and sequence analysis using emm-gene specific PCR as previously described (Beall et al., J. Clin. Micro., 1996, 34, 953-958; and Facklam et al., Emerg. Infect. Dis., 1999, 5, 247-253). Homology searches on DNA sequences were conducted against known emm sequences present in (www.cdc.gov/ncidod/biotech/infotech_hp.html). For MLST analysis, internal fragments of seven housekeeping genes, were amplified by PCR and analyzed as previously described (Enright et al., Infection and Immunity 2001, 69, 2416-2427). The emm-type was determined from comparison to the MLST database.

Broad Range Survey/Drill-Down Process (100): For *Streptococcus pyogenes*, the objective was the identification of a signature of the virulent epidemic strain and determination of its emm-type. Emm-type information is useful both for treatment considerations and epidemic surveillance. A total of 51 throat swabs were taken both from healthy recruits and from hospitalized patients in December 2002, during the peak of a GAS outbreak at a military training

camp. Twenty-seven additional isolates from previous infections ascribed to GAS were also examined. Initially, isolated colonies were examined both from throat culture samples and throat swabs directly without the culture step. The latter path can be completed within 6-12 hours providing information on a significant number of samples rapidly enough to be useful in
5 managing an ongoing epidemic.

The process of broad range survey/drill-down (200) is shown in Figure 22. A clinical sample such as a throat swab is first obtained from an individual (201). Broad range survey primers are used to obtain amplification products from the clinical sample (202) which are analyzed to determine a BCS (203) from which a species is identified (204). Drill-down primers
10 are then employed to obtain PCR products (205) from which specific information is obtained about the species (such as Emm-type) (206).

Broad Range Survey Priming: Genomic regions targeted by the broad range survey primers were selected for their ability to allow amplification of virtually all known species of bacteria and for their capability to distinguish bacterial species from each other by base
15 composition analysis. Initially, four broad-range PCR target sites were selected and the primers were synthesized and tested. The targets included universally conserved regions of 16S and 23S rRNA, and the gene encoding ribosomal protein L3 (rpoC).

While there was no special consideration of *Streptococcus pyogenes* in the selection of the broad range survey primers (which were optimized for distinguishing all important pathogens
20 from each other), analysis of genomic sequences showed that the base compositions of these regions distinguished *Streptococcus pyogenes* from other respiratory pathogens and normal flora, including closely related species of *streptococci*, *staphylococci*, and *bacilli* (Figure 23).

Drill Down Priming (Emm-Typing): In order to obtain strain-specific information about the epidemic, a strategy was designed to measure the base compositions of a set of fast clock
25 target genes to generate strain-specific signatures and simultaneously correlate with emm-types. In classic MLST analysis, internal fragments of seven housekeeping genes (*gki*, *gtr*, *murI*, *mutS*, *recP*, *xpt*, *yqiL*) are amplified, sequenced and compared to a database of previously studied isolates whose emm-types have been determined (Horner et al. Fundamental and Applied Toxicology, 1997, 36, 147). Since the analysis enabled by the present embodiment of the present
30 invention provides base composition data rather than sequence data, the challenge was to identify the target regions that provide the highest resolution of species and least ambiguous emm-classification. The data set from Table 2 of Enright et al. (Enright et al. Infection and Immunity, 2001, 69, 2416-2427) to bioinformatically construct an alignment of concatenated alleles of the seven housekeeping genes from each of 212 previously emm-typed strains, of which 101 were

unique sequences that represented 75 distinct emm-types. This alignment was then analyzed to determine the number and location of the optimal primer pairs that would maximize strain discrimination strictly on base composition data.

An example of assignment of BCSs of PCR products is shown in Figure 24 where PCR products obtained using the *gtr* primer (a drill-down emm-typing primer) from two different swab samples were analyzed (sample 12 – top and sample 10 – bottom). The deconvoluted ESI-FCTIR spectra provide accurate mass measurements of both strands of the PCR products, from which a series of candidate BCSs were calculated from the measured mass (and within the measured mass uncertainty). The identification of complementary candidate BCSs from each strand provides a means for unambiguous assignment of the BCS of the PCR product. BCSs and molecular masses for each strand of the PCR product from the two different samples are also shown in Figure 24. In this case, the determination of BCSs for the two samples resulted in the identification of the emm-type of *Streptococcus pyogenes* – sample 12 was identified as emm-type 3 and sample 10 was identified as emm-type 6.

The results of the composition analysis using the six primer pairs, 5'-emm gene sequencing and MLST gene sequencing method for the GAS epidemic at a military training facility are compared in Figure 25. The base composition results for the six primer pairs showed a perfect concordance with 5'-emm gene sequencing and MLST sequencing methods. Of the 51 samples taken during the peak of the epidemic, all but three had identical compositions and corresponded to emm-type 3. The three outliers, all from healthy individuals, probably represent non-epidemic strains harbored by asymptomatic carriers. Samples 52-80, which were archived from previous infections from Marines at other naval training facilities, showed a much greater heterogeneity of composition signatures and emm-types.

Example 19: Base Composition Probability Clouds

Figure 18 illustrates the concept of base composition probability clouds via a pseudo-four dimensional plot of base compositions of enterobacteria including *Y. pestis*, *Y. pseudotuberculosis*, *S. typhimurium*, *S. typhi*, *Y. enterocolitica*, *E. coli* K12, and *E. coli* O157:H7. In the plot of Figure 18, A, C and G compositions correspond to the x, y and z axes respectively whereas T compositions are represented by the size of the sphere at the junction of the x, y and z coordinates. There is no absolute requirement for having a particular nucleobase composition associated with a particular axis. For example, a plot could be designed wherein G, T and C compositions correspond to the x, y and z axes respectively whereas the A composition corresponds to the size of the sphere at the junction of the x, y and z coordinates. Furthermore, a

different representation can be made of the "pseudo fourth" dimension i.e.: other than the size of the sphere at junction of the x, y and z coordinates. For example, a symbol having vector information such as an arrow or a cone can be rotated at an angle that varies proportionally with the composition of the nucleobase corresponding to the pseudo fourth dimension. The choice of
5 axes and pseudo fourth dimensional representation is typically made with the aim of optimal visualization of the data being presented.

A similar base composition probability cloud analysis has been presented for a series of viruses in U.S. provisional patent application Serial No. 60/431,319, which is commonly owned and incorporated herein by reference in its entirety. In this base composition probability cloud
10 analysis, the closely related Dengue virus types 1-4 are clearly distinguishable from each other. This example is indicative of a challenging scenario for species identification based on BCS analysis because RNA viruses have a high mutation rate, it would be expected to be difficult to resolve closely related species. However, as this example illustrates, BCS analysis, aided by base composition probability cloud analysis is capable of resolution of closely related viral
15 species.

A base composition probability cloud can also be represented as a three dimensional plot instead of a pseudo-four dimensional plot. An example of such a three dimensional plot is a plot of G, A and C compositions correspond to the x, y and z axes respectively, while the composition of T is left out of the plot. Another such example is a plot where the compositions
20 of all four nucleobases is included: G, A and C+T compositions correspond to the x, y and z axes respectively. As for the pseudo-four dimensional plots, the choice of axes for a three dimensional plot is typically made with the aim of optimal visualization of the data being presented.

Example 20: Biochemical Processing of Large Amplification Products for Analysis by Mass 25 Spectrometry

In the example illustrated in Figure 26, a primer pair which amplifies a 986 bp region of the 16S ribosomal gene in *E. coli* (K12) was digested with a mixture of 4 restriction enzymes: *Bst*NI, *Bsm*FI, *Bfa*I, and *Nco*I. Figure 26(a) illustrates the complexity of the resulting ESI-FTICR mass spectrum that contains multiple charge states of multiple restriction fragments.
30 Upon mass deconvolution to neutral mass, the spectrum is significantly simplified and discrete oligonucleotide pairs are evident (Figure 26b). When base compositions are derived from the masses of the restriction fragments, perfect agreement is observed for the known sequence of nucleotides 1-856 (Figure 26c); the batch of *Nco*I enzyme used in this experiment was inactive and resulted in a missed cleavage site and a 197-mer fragment went undetected as it is outside

the mass range of the mass spectrometer under the conditions employed. Interestingly however, both a forward and reverse strand were detected for each fragment measured (solid and dotted lines in, respectively) within 2 ppm of the predicted molecular weights resulting in unambiguous determination of the base composition of 788 nucleotides of the 985 nucleotides in the amplicon.

- 5 The coverage map offers redundant coverage as both 5' to 3' and 3' to 5' fragments are detected for fragments covering the first 856 nucleotides of the amplicon.

This approach is in many ways analogous to those widely used in MS-based proteomics studies in which large intact proteins are digested with trypsin, or other proteolytic enzyme(s), and the identity of the protein is derived by comparing the measured masses of the tryptic
10 peptides with theoretical digests. A unique feature of this approach is that the precise mass measurements of the complementary strands of each digest product allow one to derive a de novo base composition for each fragment, which can in turn be "stitched together" to derive a complete base composition for the larger amplicon. An important distinction between this approach and a gel-based restriction mapping strategy is that, in addition to determination of the
15 length of each fragment, an unambiguous base composition of each restriction fragment is derived. Thus, a single base substitution within a fragment (which would not be resolved on a gel) is readily observed using this approach. Because this study was performed on a 7 Tesla ESI-FTICR mass spectrometer, better than 2 ppm mass measurement accuracy was obtained for all fragments. Interestingly, calculation of the mass measurement accuracy required to derive
20 unambiguous base compositions from the complementary fragments indicates that the highest mass measurement accuracy actually required is only 15 ppm for the 139 bp fragment (nucleotides 525-663). Most of the fragments were in the 50-70 bp size-range which would require mass accuracy of only ~50 ppm for unambiguous base composition determination. This level of performance is achievable on other more compact, less expensive MS platforms such as
25 the ESI-TOF suggesting that the methods developed here could be widely deployed in a variety of diagnostic and human forensic arenas.

This example illustrates an alternative approach to derive base compositions from larger PCR products. Because the amplicons of interest cover many strain variants, for some of which complete sequences are not known, each amplicon can be digested under several different
30 enzymatic conditions to ensure that a diagnostically informative region of the amplicon is not obscured by a "blind spot" which arises from a mutation in a restriction site. The extent of redundancy required to confidently map the base composition of amplicons from different markers, and determine which set of restriction enzymes should be employed and how they are most effectively used as mixtures can be determined. These parameters will be dictated by the

extent to which the area of interest is conserved across the amplified region, the compatibility of the various restriction enzymes with respect to digestion protocol (buffer, temperature, time) and the degree of coverage required to discriminate one amplicon from another.

Example 21: Identification of members of the Viral Genus *Orthopoxvirus*

- 5 Primer sites were identified on three essential viral genes – the DNA-dependent polymerase (DdDp), and two sub-units of DNA-dependent RNA polymerases A and B (DdRpA and DdRpB). These intelligent primers designed to identify members of the viral genus *Orthopoxvirus* are shown in Table 12 wherein Tp = 5' propynylated uridine and Cp = 5' propynylated cytidine.

10

Table 12

**Intelligent Primer Pairs for Identification of members of the
Viral Genus *Orthopoxvirus***

Primer Pair Name	Forward Primer Sequence	Forward SEQ ID NO:	Reverse Primer Sequence	Reverse SEQ ID NO:
A25L NC00161 1 28 127	GTACTGAATCCGCCTAAG	354	GTGAATAAAGTATCGCCCTAA TA	355
A18R NC00161 1 100 207	GAAGTTGAACCGGGATCA	356	ATTATCGGTCGTTGTTAATGT	357
A18R NC00161 1 1348 1445	CTGTCTGTAGATAAACTAGGAT T	358	CGTTCTTCTCTGGAGGAT	359
E9L NC001611 1119 1222	CGATACTACGGACGC	360	CTTTATGAATTACTTTACATA T	361
K8R NC001611 221 311	CTCCTCCATCACTAGGAA	362	CTATAACATTCAAAGCTTATT G	363
A24R NC00161 1 795 878	CGCGATAATAGATAGTGCTAAA C	364	GCTTCCACCAGGTCATTAA	365
A25L NC00161 1 28 127P	GTACpTpGAATpCpCpGCpCpT AAG	366	GTGAATAAAGTATpCpGCpCp CpTpAATA	367
A18R NC00161 1 100 207P	GAAGTpTpGAACpCpGGGATCA	368	ATTATCGGTpCpGTpTpGTpT pAATGT	369
A18R NC00161 1 1348 1445P	CTGTpCpTpGTAGATAAACpTp AGGATT	370	CGTTCpTpTpCpTpCpTpGGA GGAT	371
E9L NC001611 1119 1222P	CGATACpTpACpGGACGC	372	CTTTATGAATpTpACpTpTpT pACATAT	373
K8R NC001611 221 311P	CTpCpCpTCpCpATCACpTpAG GAA	374	CTATAACATpTpCpAAAGCpT pTpATTG	375
A24R NC00161 1 795 878P	CGCGATpAATpAGATAGTpGCp TpAAAC	376	GCTTCpCpACpCAGGTpCATp TAA	377

- As illustrated in Figure 27, members of the *Orthopoxvirus* genus group can be identified, distinguished from one another, and distinguished from other members of the Poxvirus family using a single pair of primers designed against the DdRpB gene.

Since the primers were designed across regions of high conservation within this genus, the likelihood of missed detection due to sequence variations at these sites is minimized. Further,

none of the primers is expected to amplify other viruses or any other DNA, based on the data available in GenBank. This method can be used for all families of viral threat agents and is not limited to members of the *Orthopoxvirus* genus.

5 Example 22: Identification of Viruses that Cause Viral Hemorrhagic Fevers

In accordance with the present invention an approach of broad PCR priming across several different viral species is employed using conserved regions in the various viral genomes, amplifying a small, yet highly informative region in these organisms, and then analyzing the resultant amplicons with mass spectrometry and data analysis. These regions will be tested with
10 live agents, or with genomic constructs thereof.

Detection of RNA viruses will necessitate a reverse transcription (RT) step prior to the PCR amplification of the TIGER reporter amplicon. To maximize throughput and yield while minimizing the handling of the samples, commercial one-step reverse transcription polymerase chain reaction (RT-PCR) kits will be evaluated for use. If necessary, a one-step RT-PCR mix
15 using our selected DNA polymerase for the PCR portion of the reaction will be developed. To assure there is no variation in our reagent performance all new lots of enzymes, nucleotides and buffers will be individually tested prior to use.

Various modifications of the invention, in addition to those described herein, will be apparent to those skilled in the art from the foregoing description. Such modifications are also
20 intended to fall within the scope of the appended claims. Each reference, web site, Genebank accession number, etc. cited in the present application is incorporated herein by reference in its entirety. The following US applications are incorporated herein by reference in their entirety: Serial No. 10/323,233 filed December 18, 2002, Serial No. 10/326,051 filed December 18, 2002, Serial No. 10/325,527 filed December 18, 2002, Serial No. 10/325,526 filed December 18, 2002,
25 Serial No. 60/431,319 filed December 6, 2002, Serial No. 60/443,443 filed January 29, 2003, Serial No. 60/443,788 filed January 30, 2003, Serial No. 60/447,529 filed February 14, 2003, and Serial No. 60/501,926 filed September 11, 2003.

What is claimed is:

1. A method of identifying a plurality of etiologic agents of disease in an individual comprising the steps of:
amplifying at least one nucleic acid molecule obtained from a biological sample from
5 the individual with a plurality of intelligent primers to obtain a plurality of amplification products corresponding to the plurality of etiologic agents; and
determining the molecular masses of the plurality of amplification products, wherein the molecular masses identify the plurality of etiologic agents and wherein the intelligent primers are broad range survey primers, division-wide primers, drill-down primers, or any combination
10 thereof.
2. A method of claim 1 wherein identification of at least one of the plurality of etiologic agents is accomplished at the genus or species level, and the intelligent primers are broad range survey primers, division-wide primers, or any combination thereof.
3. A method of claim 1 wherein a subspecies characteristic of at least one of the plurality
15 of etiologic agents is obtained using drill-down primers.
4. A method of claim 3 wherein the subspecies characteristic is serotype, strain type, sub-strain type, sub-species type, emm-type, presence of a bioengineered gene, presence of a toxin gene, presence of an antibiotic resistance gene, presence of a pathogenicity island, or presence of a virulence factor.
- 20 5. A method of claim 1 wherein the molecular mass is determined by mass spectrometry.
6. A method of claim 5 wherein the mass spectrometry is Fourier transform ion cyclotron resonance mass spectrometry (FTICR- MS), ion trap, quadrupole, magnetic sector, time of flight (TOF), Q-TOF, or triple quadrupole.
7. A method of claim 1 wherein the molecular masses are used to determine the base
25 compositions of the amplification products and wherein the base compositions identify the pathogen.
8. A method of *in silico* screening of intelligent primer sets for identification of a plurality of bioagents comprising the steps of:
preparing a base composition probability cloud plot from a plurality of base
30 composition signatures of the plurality of bioagents generated *in silico*;
inspecting the base composition probability cloud plot for overlap of clouds from different bioagents; and
selecting primer sets based on minimal overlap of the clouds.
9. A method of performing epidemic surveillance comprising the steps of:

determining the molecular mass of the amplification product, wherein said molecular mass identifies the pathogen in the biological sample.

28. A method of claim 27 wherein the pathogen is a bacterium, a virus, a protozoan, a parasite, a mold, or a fungus.
- 5 29. A method of claim 27 wherein the biological sample is blood, mucus, hair, urine, breath, sputum, saliva, stool, nail, or tissue biopsy.
30. A method of claim 27 wherein the biological sample is obtained from an animal.
31. A method of claim 30 wherein the animal is a human.
32. A method of claim 27 wherein the intelligent primers are broad range survey primers,
10 division-wide primers, or drill-down primers.
33. A method of claim 32 wherein identification of the pathogen is accomplished at the genus or species level, and wherein the intelligent primers are broad range survey primers or division-wide primers.
34. A method of claim 32 wherein a subspecies characteristic of the pathogen is obtained
15 using drill-down primers.
35. A method of claim 34 wherein the subspecies characteristic is serotype, strain type, sub-strain type, sub-species type, emm-type, presence of a bioengineered gene, presence of a toxin gene, presence of an antibiotic resistance gene, presence of a pathogenicity island, or presence of a virulence factor.
- 20 36. A method of claim 27 wherein the molecular mass is determined by mass spectrometry.
37. A method of claim 36 wherein the mass spectrometry is Fourier transform ion cyclotron resonance mass spectrometry (FTICR- MS), ion trap, quadrupole, magnetic sector, time of flight (TOF), Q-TOF, or triple quadrupole.
38. A method of claim 27 wherein the intelligent primers are targeted to ribosomal RNA or
25 housekeeping genes.
39. A method of claim 27 wherein the molecular mass is used to determine the base composition of said amplification product and wherein said base composition identifies said pathogen.
40. An intelligent primer pair wherein each member of the pair has at least 70% sequence
30 identity with the sequence of the corresponding member of any one of the following intelligent primer pair sequences: SEQ ID NOs: 8:9, 10:11, 12:13, 14:15, 16:17, 18:19, 20:21, 22:23, 24:25, 26:27, 28:29, 30:31, 32:33, 34:35, 36:37, 38:39, 40:41, 42:43, 44:45, 46:47, 48:49, 50:51, 52:53, 54:55, 56:57, 58:59, 60:61, 62:63, 64:65, 66:67, 68:69, 70:71, 72:73, 74:75, 76:77, 78:79, 80:81, 82:83, 84:85, 86:87, 88:89, 90:91, 92:93, 94:95, 96:97, 98:99, 100:101, 102:103, 104:105,

106:107, 108:109, 110:111, 112:113, 114:115, 116:117, 118:119, 120:121, 122:123, 124:125,
126:127, 128:129, 130:131, 132:133, 134:135, 136:137, 138:139, 140:141, 142:143, 144:145,
146:147, 148:149, 150:151, 152:153, 154:155, 156:157, 158:159, 160:161, 162:163, 164:165,
166:167, 168:169, 170:171, 172:173, 174:175, 176:177, 178:179, 180:181, 182:183, 184:185,
5 186:187, 188:189, 190:191, 192:193, 194:195, 196:197, 198:199, 200:201, 202:203, 204:205,
206:207, 208:209, 210:211, 212:213, 214:215, 216:217, 218:219, 220:221, 222:223, 224:225,
226:227, 228:229, 230:231, 232:233, 234:235, 236:237, 238:239, 240:241, 242:243, 244:245,
246:247, 248:249, 250:251, 252:253, 254:255, 256:257, 258:259, 260:261, 262:263, 264:265,
266:267, 268:269, 270:271, 272:273, 274:275, 276:277, 278:279, 280:281, 282:283, 284:285,
10 286:287, 288:289, 290:291, 292:293, 294:295, 296:297, 298:299, 300:301, 302:303, 304:305,
306:307, 308:309, 310:311, 312:313, 314:315, 316:317, 318:319, 320:321, 322:323, 324:325,
326:327, 328:329, 330:331, 332:333, 334:335, 336:337, 338:339, 340:341, 342:343, 344:345,
346:347, 348:349, 350:351, 352:353, 354:355, 356:357, 358:359, 360:361, 362:363, 364:365,
366:367, 368:369, 370:371, 372:373, 374:375, or 376:377.

15 41. The intelligent primer pair of claim 40 comprising at least one modified nucleobase.

42. The intelligent primer pair of claim 41 wherein the modified nucleobase is 5-propynylcytidine or 5-propynyluridine.

43. A bioagent identifying amplicon comprising an isolated polynucleotide of about 45 to
about 150 nucleobases in length produced by the process of amplification of nucleic acid from a
20 bioagent with a pair of intelligent primers wherein each intelligent primer is of a length of about
12 to about 35 nucleobases, wherein the bioagent identifying amplicon provides identifying
information about the bioagent.

44. The bioagent identifying amplicon of claim 43 wherein each member of the pair has at
least 70% sequence identity with the sequence of the corresponding member of any one of the
25 following intelligent primer pair sequences: SEQ ID NOs: 8:9, 10:11, 12:13, 14:15, 16:17, 18:19,
20:21, 22:23, 24:25, 26:27, 28:29, 30:31, 32:33, 34:35, 36:37, 38:39, 40:41, 42:43, 44:45, 46:47,
48:49, 50:51, 52:53, 54:55, 56:57, 58:59, 60:61, 62:63, 64:65, 66:67, 68:69, 70:71, 72:73, 74:75,
76:77, 78:79, 80:81, 82:83, 84:85, 86:87, 88:89, 90:91, 92:93, 94:95, 96:97, 98:99, 100:101,
102:103, 104:105, 106:107, 108:109, 110:111, 112:113, 114:115, 116:117, 118:119, 120:121,
30 122:123, 124:125, 126:127, 128:129, 130:131, 132:133, 134:135, 136:137, 138:139, 140:141,
142:143, 144:145, 146:147, 148:149, 150:151, 152:153, 154:155, 156:157, 158:159, 160:161,
162:163, 164:165, 166:167, 168:169, 170:171, 172:173, 174:175, 176:177, 178:179, 180:181,
182:183, 184:185, 186:187, 188:189, 190:191, 192:193, 194:195, 196:197, 198:199, 200:201,
202:203, 204:205, 206:207, 208:209, 210:211, 212:213, 214:215, 216:217, 218:219, 220:221,

222:223, 224:225, 226:227, 228:229, 230:231, 232:233, 234:235, 236:237, 238:239, 240:241, 242:243, 244:245, 246:247, 248:249, 250:251, 252:253, 254:255, 256:257, 258:259, 260:261, 262:263, 264:265, 266:267, 268:269, 270:271, 272:273, 274:275, 276:277, 278:279, 280:281, 282:283, 284:285, 286:287, 288:289, 290:291, 292:293, 294:295, 296:297, 298:299, 300:301, 5 302:303, 304:305, 306:307, 308:309, 310:311, 312:313, 314:315, 316:317, 318:319, 320:321, 322:323, 324:325, 326:327, 328:329, 330:331, 332:333, 334:335, 336:337, 338:339, 340:341, 342:343, 344:345, 346:347, 348:349, 350:351, 352:353, 354:355, 356:357, 358:359, 360:361, 362:363, 364:365, 366:367, 368:369, 370:371, 372:373, 374:375, or 376:377.

45. A bioagent identifying amplicon for identification of a bacterium comprising an isolated
10 polynucleotide of about 45 to about 150 nucleobases in length produced by the process of amplification of nucleic acid encoding ribosomal RNA from a bacterium with a pair of intelligent primers wherein each intelligent primer is of a length of about 12 to about 35 nucleobases, wherein the bioagent identifying amplicon provides identifying information about the bioagent.


15 46. The bioagent identifying amplicon of claim 45 wherein each member of the pair has at least 70% sequence identity with the sequence of the corresponding member of any one of the following intelligent primer pair sequences: SEQ ID NOs: 8:9, 10:11, 12:13, 14:15, 16:17, 18:19, 20:21, 22:23, 24:25, 26:27, 28:29, 30:31, 32:33, 34:35, 36:37, 38:39, 40:41, 42:43, 44:45, 46:47, 48:49, 50:51, 52:53, 54:55, 56:57, 58:59, 60:61, 62:63, 64:65, 66:67, 68:69, 70:71, 72:73, 74:75, 20 76:77, 78:79, 80:81, 82:83, 84:85, 86:87, 88:89, 90:91, 92:93, 94:95, 96:97, 98:99, 100:101, 102:103, 104:105, 106:107, 108:109, 110:111, 112:113, 114:115, 116:117, 118:119, 120:121, 122:123, 124:125, 126:127, 128:129, 130:131, 132:133, 134:135, 136:137, 138:139, 140:141, 142:143, 144:145, 146:147, 148:149, 150:151, 152:153, 154:155, 156:157, 158:159, 160:161, 162:163, 164:165, 166:167, 168:169, 170:171, 172:173, 174:175, 176:177, 178:179, 180:181, 25 182:183, 184:185, 186:187, 188:189, 190:191, 192:193, 194:195, 196:197, 198:199, 200:201, 202:203, 204:205, 206:207, 208:209, 210:211, 212:213, 214:215, 216:217, 218:219, 220:221, 222:223, 224:225, 226:227, 228:229, 230:231, 232:233, 234:235, 236:237, 238:239, 240:241, 242:243, 244:245, 246:247, 248:249, 250:251, 252:253, 254:255, 256:257, 258:259, 260:261, 262:263, 264:265, 266:267, 268:269, 270:271, 272:273, 274:275, 276:277, 278:279, 280:281, 30 282:283, 284:285, 334:335, 336:337, 338:339, 340:341, 342:343, 344:345, 346:347, 348:349, 350:351, or 352:353.

47. A bioagent identifying amplicon for identification of a virus comprising an isolated polynucleotide of about 45 to about 150 nucleobases in length produced by the process of amplification of nucleic acid encoding a viral housekeeping gene with a pair of intelligent

primers wherein each intelligent primer is of a length of about 12 to about 35 nucleobases, wherein the bioagent identifying amplicon provides identifying information about the bioagent.

48. The bioagent identifying amplicon of claim 47 wherein each member of the pair has at least 70% sequence identity with the sequence of the corresponding member of any one of the
- 5 following intelligent primer pair sequences: SEQ ID NOs: 286:287, 288:289, 290:291, 292:293, 294:295, 296:297, 298:299, 300:301, 302:303, 304:305, 306:307, 308:309, 310:311, 312:313, 314:315, 316:317, 318:319, 320:321, 322:323, 324:325, 326:327, 328:329, 330:331, 332:333, 354:355, 356:357, 358:359, 360:361, 362:363, 364:365, 366:367, 368:369, 370:371, 372:373, 374:375, or 376:377.
- 10 49. The method of claim 48 wherein said viral housekeeping gene is hexon, DNA-dependent polymerase, DNA-dependent RNA polymerase A, or DNA-dependent RNA polymerase B.

FIG. 1A-1






FIG. 1A-3





FIG. 1A-4







FIG. 1C






FIG. 1E




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7





FIG. 8

ESI-TOF MS of sspE 56mer + Calibrant






FIG. 10





FIG. 11







FIG. 14



Picorno RdRp 705-759

■ Encephalomyocarditis virus	□ Foot-and-mouth disease virus	▨ Polio	□ Rhinovirus
▨ Enterovirus	■ Hepatitis A virus	■ Porcine enterovirus	▨ Simian Hepatitis A

FIG. 15





FIG. 16



Flavi RdRp 2453-2493

Dengue virus type
 Dengue virus type
 Dengue virus type

Japanese encephalitis virus
 Kunjin virus
 Murray valley encephalitis virus

Tick-borne encephalitis virus
 West Nile virus
 Yellow fever virus

FIG. 17




Figure 18





Figure 19






Figure 20

Figure 21




Figure 22

Universal Survey/Drill-Down Process




Figure 23

Base Composition Signatures from primer pair 14 (16S rRNA)




Figure 24




Figure 25

	EMM-type determination			Base Compositions										
	Sample	MLST- <i>emm1</i>	Reagendin- <i>emr</i>	<i>murt</i>	<i>mutS</i>	<i>mutL</i>	<i>vall</i>	<i>akl</i>	<i>gtr</i>					
Outbreak Samples	1	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32					
	2	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32					
	3	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32					
	4	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32					
	5	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32					
	6	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32					
	7	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32					
	8	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32					
	9	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32					
	10	6	6	A40 G24 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A31 G35 C17 T32	A39 G28 C16 T32					
	11	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32					
	12	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32					
	13	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32					
	14	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32					
	15	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32					
	16	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32					
	17	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32					
	18	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32					
	19	6	6	A40 G24 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A31 G35 C17 T32	A39 G28 C16 T32					
	20	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32					
	21	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32					
	22	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32					
	23	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32					
	24	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32					
	25	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32					
	26	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32					
27	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
28	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
29	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
30	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
31	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
32	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
33	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
34	28	28	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
35	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
36	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
37	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
38	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
39	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
40	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
41	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
42	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
43	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
44	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
45	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
46	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
47	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
48	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
49	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
50	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
51	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
52	44/61/82.0	44/61	A40 G24 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
53	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
54	44/61/82.0	44/61	A40 G24 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
55	11	11	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
56	25.75	75	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
57	6	6	A40 G24 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A31 G35 C17 T32	A39 G28 C16 T32						
58	25.75	75	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
59	12	12	A40 G24 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A31 G35 C17 T32	A39 G28 C16 T32						
60	25.75	75	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
61	6	6	A40 G24 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A31 G35 C17 T32	A39 G28 C16 T32						
62	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
63	6	6	A40 G24 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A31 G35 C17 T32	A39 G28 C16 T32						
64	5.59	5	A40 G24 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
65	8	5	A40 G24 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
66	5.59	5	A40 G24 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
67	8	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
68	44/61/82.0	44/61	A40 G24 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
69	5.59	5	A40 G24 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
70	3	3	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
71	53.91	29	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
72	44/61/82.0	44/61	A40 G24 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
73	6	6	A40 G24 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A31 G35 C17 T32	A39 G28 C16 T32						
74	6	6	A40 G24 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A31 G35 C17 T32	A39 G28 C16 T32						
75	12	12	A40 G24 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A31 G35 C17 T32	A39 G28 C16 T32						
76	44/61/82.0	44/61	A40 G24 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
77	3	3 [50% identity]	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
78	12	12	A40 G24 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A31 G35 C17 T32	A39 G28 C16 T32						
79	22	3 [50% identity]	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						
80	53.91	29	A39 G25 C20 T34	A38 G27 C23 T33	A30 G36 C20 T36	A40 G29 C19 T31	A32 G35 C17 T32	A39 G28 C16 T32						

Figure 26





Figure 27



SEQUENCE LISTING

<110> ISIS Pharmaceuticals, Inc.

Ecker, David J.

Griffey, Richard H.

Sampath, Rangarajan

Hofstadler, Steven

McNeil, John

Crooke, Stanley T.

Blyn, Lawrence B.

Ranken, Raymond

Hall, Thomas A.

<120> METHODS FOR RAPID IDENTIFICATION OF PATHOGENS IN HUMANS AND ANIMALS

<130> IBIS0060-500WO (DIBIS-0044WO)

<150> 60/431,319

<151> 2002-12-06

<150> 10/323,233

<151> 2002-12-18

<150> 10/326,051

<151> 2002-12-18

<150> 10/325,527

<151> 2002-12-18

<150> 10/325,526

<151> 2002-12-18

<150> 60/443,443

<151> 2003-01-29

<150> 60/443,788

<151> 2003-01-30

<150> 60/447,529

<151> 2003-02-14

<150> 60/501,926

<151> 2003-09-11

<160> 377

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 1388

<212> RNA

<213> 16S rRNA Consensus Sequence

<220>

<221> misc_feature

<222> 1-7, 15, 22-24, 36-38, 41, 42,44, 56, 59-90, 93, 97, 98, 109,
110, 112-116, 118-120, 123-131, 134,136, 138-144, 149-155, 161,
162, 164-177, 182-209, 212-220, 222-225, 227, 230, 231, 235, 236,
241, 245, 246, 253, 255-257, 260, 261-263, 267, 269, 270, 278, 279,

281, 282, 284, 291, 294, 301, 306, 310, 329, 330, 335, 336-338,
 344, 345, 347, 350, 351, 355, 356, 357, 361, 363, 364, 371, 372,
 373, 376, 379, 382-386, 388, 389, 394-396, 398-405, 408, 411-438,
 442, 443, 445-451, 453, 454, 458-460, 465, 469, 491, 495, 496, 499,
 504-506, 511, 512, 514, 524, 526-528, 530, 534, 537-544, 546-550,
 552, 556-562, 565, 569-578, 580-586, 589-595, 597, 601-606, 609,
 612-617, 621-624, 629, 633, 636, 639, 643, 645, 646, 648, 650, 654,
 658-660, 669-674, 678-683, 689, 691, 693-696, 704, 708, 713, 734,
 737, 738, 744, 746-754, 756-758, 760-782, 784-786, 791-793,
 796-800, 815, 816, 823-825, 834, 845, 848, 857, 859, 864, 869, 875,
 877, 878, 884, 886, 894-898, 903-917, 921, 922-948, 955, 961, 972,
 973, 978, 990, 1005-1013, 1015, 1017, 1019, 1021-1029, 1031, 1033,
 1037-1043, 1049-1051, 1053, 1054, 1057-1059, 1069, 1075, 1083, 1085,
 1089, 1094, 1096-1099, 1104, 1110, 1111, 1119-1123, 1127, 1128,
 1130, 1132, 1133, 1136, 1138, 1139-1141, 1143, 1144, 1146-1150,
 1154, 1157, 1159-1162, 1166-1170, 1172, 1173, 1176, 1181, 1183-1186,
 1195-1198, 1200, 1205, 1206, 1210, 1220-1222, 1227, 1229, 1231-1233,
 1244, 1249, 1266-1268, 1271, 1273, 1274, 1277-1285, 1288, 1289,
 1293-1304, 1306-1311, 1313-1322, 1324, 1326, 1329, 1330-1338, 1341,
 1345-1347, 1361, 1364, 1366-1368, 1372, 1373, 1375, 1378, 1388

<223> n = A,T,C or G

<400> 1

```

nnnnnnnnnaga ggacnggccca gnnngaacgc ggcggnnngc nnanacagca agcgancggn 60
nnnnnnnnnnnn nnnnnnnnnnnn nnnnnnnnnnnn agngggcnaac ggaggagaann cnnnnnnnnnn 120
ccnnnnnnnnnn nggnanannnn nnnnggaaann nnnnnnaaacc nnnnnnnnnnnn nnnnnnnnaaa 180
gnnnnnnnnnnn nnnnnnnnnnnn nnnnnnnnnng annnnnnnnnn gnnnnnanagn ngggngggaa 240
nggcnnacca agncnnngan nnnagcngnn cgagaggngg nncngccaca nggnacgaga 300
nacgggccan acccacggga ggcagcagnn ggaannnnca aggnngnaan ncgannnagc 360
nannccgcgg nnnngangang gnnnnngnng aaannncnnn nnnnnnganga nnnnnnnnnnn 420
nnnnnnnnnnnn nnnnnnnnnnga cnnannnnnnn nannaagnnn cggcnaacnc ggccagcagc 480
cgcggaacag naggnngcna gcgnnnccga nnnngggcga aagngnnngn agngggnnnn 540
nnnnngnnnnn gnaaannnnnn nngcnaacnn nnnnnnnncn nnnnnnacnn nnnnnncngag 600
nnnnnnnagng gnnnnnnngaa nnnnggagng ggnaancgna ganannngnan gaanaccnnn 660
gcgaaggcnn nnnnccggnnn nnnacgacnc nnnnnncgaa agcngggngag cnaacaggaa 720
gaacccggag ccangcnaaa acngnnnnnnn nnnngnnngn nnnnnnnnnnnn nnnnnnnnnnn 780
nnaannnaacg nnaaannnnnn ccgccgggga gacgnncgca agnnnaaacc aaangaagac 840
ggggnccngc acaagcngng gagnaggga acgangnnac gcgnanaacc accnnnnnga 900
cannnnnnnnn nnnnnnnngan annnnnnnnnn nnnnnnnnnn nnnnnnnnnac agggngcagg 960
ngcgagccg gnnnggagng ggaagcccg aacgagcgca accnnnnnnn nnnngcncanc 1020
nnnnnnnnng ngnaccnnnn nnnacgccnn ngnaannng gaggaaggng gggangacgc 1080
aancncagnc ccangnnnnng ggcncacacn ncacaaggnn nnnacanngn gnnngcnnnn 1140
ngnnannnnn agcnaancnn nnaaannnnn cnnagncgga ngnnnnccgca accgnnnnncn 1200
gaagnggan cgagaacgn nnacagnang nnnccgggaa cgncgggnc gacacaccgc 1260
ccgcannnca ngnnagnnnn nnnnnccnna agnnnnnnnnn nnnnnnnnnn ngnnnnnnnn 1320
nncnanggnn nnnnnnnnga ngggnnnaag cgaacaagga nccnannnga annngngngg 1380
acaccccn

```

<210> 2

<211> 2654

<212> RNA

<213> Artificial Sequence

<220>

<221> misc_feature

<222> 1-4, 8-12, 16, 18-22, 31, 35-40, 43, 47, 54, 56-62, 64, 65, 67-69, 71, 72, 74-76, 79, 80, 83, 84, 86-93, 95-99, 101, 104-106, 108, 109, 119, 125-142, 144-171, 173-175, 178-182, 186, 194, 198-202, 208, 214, 215, 218-221, 224, 226-229, 233, 244-246, 248, 250, 251, 254, 256-280, 282-284, 288-292, 295, 299-307, 309-311, 314, 316, 318, 319, 322, 325-328, 332-354, 358-360, 362-365, 367, 370, 372, 373, 375-378, 380, 381, 385-387, 389-392, 398-403, 407, 414-420, 425, 429, 431, 433, 434, 439, 443, 451, 458, 463, 464, 465, 466, 467, 469, 479, 482, 483, 492-496, 498, 500, 503-505, 507-509, 512-525, 528, 529, 533, 537, 539, 540, 543-545, 547, 549, 552-561, 564, 567, 570-578, 580, 583, 586, 589, 594, 599-601, 604, 605, 607, 611-613, 616-620, 622-625, 630, 635, 637-639, 643, 646-648, 651, 652, 657, 662-666, 670, 672-676, 682, 689-696, 703-708, 714, 715, 718-720, 722, 725, 730, 731, 733, 736, 738, 742-744, 746, 747, 756, 757, 763-766, 770-773, 776-791, 794, 805-814, 817-829, 832, 833, 835-842, 847, 852, 855-870, 872, 875, 876, 878, 879, 881-883, 885, 887, 889, 892-894, 896-898, 900, 901, 903, 908, 913, 920, 922, 923, 925-927, 932, 936, 939-946, 952, 956, 959, 962-967, 969, 970, 972, 976-978, 983, 999, 1001, 1002, 1008, 1009, 1015, 1022, 1023, 1025, 1028-1034, 1039, 1042, 1043, 1045, 1047, 1052, 1056-1063, 1069-1074, 1076-1097, 1102, 1103, 1109-1121, 1126-1132, 1134, 1135, 1137-1143, 1147-1155, 1159, 1161, 1164, 1165, 1167, 1168, 1170, 1174, 1178-1185, 1189, 1191, 1192, 1194-1198, 1200, 1204, 1206-1208, 1210, 1215, 1218-1223, 1225, 1227-1229, 1231-1236, 1240, 1245-1247, 1253, 1254, 1258, 1260, 1263, 1265, 1267, 1268, 1271, 1272, 1277, 1278, 1280-1282, 1285, 1286, 1291-1293, 1296-1316, 1321-1326, 1328-1345, 1348-1455, 1457, 1458, 1464-1490, 1496, 1497, 1511, 1513-1516, 1518, 1519, 1523, 1525, 1526, 1528, 1529, 1533, 1535-1537, 1539-1542, 1545-1552, 1560, 1561, 1567-1571, 1576, 1581, 1583, 1588-1591, 1593-1633, 1635-1638, 1640-1642, 1644-1646, 1648-1654, 1656, 1661, 1662, 1673, 1674, 1676, 1677, 1680, 1683, 1684, 1687, 1691, 1692, 1695, 1699, 1702, 1703, 1707, 1714, 1718, 1719, 1727, 1728, 1730-1738, 1740-1744, 1746-1756, 1758-1760, 1767, 1768, 1770, 1779, 1780, 1789, 1790, 1820, 1828, 1831, 1833, 1836, 1839-1846, 1851-1859, 1861, 1863, 1865, 1869-1871, 1873-1877, 1879, 1886-1890, 1892, 1896-1900, 1915, 1916, 1918, 1920, 1924, 1925, 1927-1932, 1934, 1936-1950, 1952, 1953, 1956, 1961, 1966, 1968, 1969, 1970, 1973-1980, 1983, 1984, 1987-1993, 1998, 2000-2004, 2006, 2007, 2011, 2014, 2016-2029, 2034-2044, 2046, 2048-2056, 2061, 2063-2065, 2067, 2068, 2072, 2075, 2085, 2086, 2091, 2095, 2096, 2106, 2108, 2109, 2111, 2116-2118, 2120, 2122-2125, 2128, 2129, 2132, 2133, 2136-2143, 2146, 2147, 2150, 2151, 2153, 2155, 2159, 2160, 2161, 2164, 2165, 2169, 2170, 2173-2176, 2179-2182, 2190-2192, 2199, 2200, 2203-2205, 2214, 2217-2222, 2228, 2232, 2248, 2251, 2253, 2266, 2268-2271, 2280, 2283, 2291-2294, 2311, 2313, 2324, 2327, 2328, 2339, 2340, 2349, 2350, 2355, 2358, 2362, 2370, 2372, 2386, 2394, 2396, 2397, 2399, 2401, 2403, 2405-2407, 2410-2412, 2415-2417, 2419-2421, 2423, 2424, 2426, 2442-2444, 2446, 2449, 2450, 2452, 2454, 2459-2461, 2463, 2466-2468, 2473-2475, 2479, 2480, 2483, 2485, 2486, 2491-2494, 2497-2500, 2505, 2506, 2512, 2520-2522, 2526, 2528-2530, 2532-2535, 2539, 2540, 2543-2545, 2547, 2549-2568, 2571-2573, 2575, 2576-2579, 2583, 2584, 2586-2589, 2591, 2597-2599, 2601-2603, 2605, 2609-2612, 2614, 2615, 2617, 2618-2622, 2625-2627, 2630-2632, 2637-2640, 2642-2647, 2649-2654

<223> n = A,T,C or G

<400> 2

nnnaagnnnn nnaagngnnnn nnggggagcc nggcnnnnnn agncgangaa ggangnnnnnn 60
nnccnnnnna nncnnngggn agngnnnnnn nnccnnnnna nccnnngnnc cgaaggggna 120

```

accnnnnnnnn nnnnnnnnnn nnannnnnnnn nnnnnnnnnn nnnnnnnnnn ngnnnacnnn 180
nngaangaaa cacnagannn nnaggaanag aaannaannn ngancnnnnn agnggcgagc 240
gaannngnan nagnnnnnnnn nnnnnnnnnn nnnnnnnnnn annngaannn nnggnaagnn 300
nnnnnnnann nggnanannc cngannnnnaa annnnnnnnn nnnnnnnnnn nnnnagannn 360
cnnnnncnngn gnnannnnngn ngaannngnn nggaccannn nnaagncaa aacnnnnnnn 420
gaccnaagng nannagacng ganggaaag ngaaaagnac cnnnnnnang ggaggaaana 480
gnnccgaaac cnnnnncnan aannngnnna gnnnnnnnnn nnnnnngann cgnccgnann 540
agnnncngng annnnnnnnn ngcnagnaann nnnnnnnngn agncgnagng aaancgagnn 600
naannngcg nnnagnnnnn gnnnnagacn cgaancnnng gancannnag nncaggngaa 660
gnnnnnngaann annnnnnggag gnccgaaacnn nnnnnnggaa aannnnnnng agannngnnn 720
gnggngaann ncnanncnaa cnnngnnaag cgcccnncga aannnnnaggn nnnngcnnnnn 780
nnnnnnnnnn nggnggagag cagcgnnnnnn nnnngggnnn nnnnnnnnna cnnannnnnn 840
nnaaacnnga accnnnnnnn nnannnnnnn gnagnnann nngngngna annncnnngn 900
nanagggnaa canccagan cnnnnnaag gncnnaann nnnnnnaagg gnaaangang 960
gnnnnnncnn anacannnag gangggcaga agcagccanc nnaaagann cgaanagcca 1020
cnnncagnnn nnnngcgcng annanancgg gncannnnn nnnccgaann nnnngnnnnn 1080
nnnnnnnnnn nnnnnnnngga gngagcggn nnnnnnnnnn ngaagnnnn nngnnannnn 1140
nnnggannnn nnnnnagng ngngnnnnn agancgannn nnnnnngana nncnnnnncn 1200
cgnannnncn aaggncennn nnnangnnnc nnnnnngggn agcgnnncca agnngagncn 1260
ganangnnag nngaggnnan nnggnnaacc nnnacnnnnn nnnnnnnnnn nnnnnngacg 1320
nnnnnnngnn nnnnnnnnnn nnnnnnggnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 1380
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 1440
nnnnnnnnnn nnnnnncnga aaannnnnnn nnnnnnnnnn nnnnnnnnnn cgaccnnaaa 1500
ccgacacagg ngnnnnngng agnanncnna gngngnnngn nnaannnnnn nnaaggaacn 1560
ngcaannnnn nccganccgg nanaagggnn ncnnnnnnnn nnnnnnnnnn nnnnnnnnnn 1620
nnnnnnnnnn nnnngnnnnn nnannngnnn nnnncnacga nnaaaaacac agnncnngcn 1680
aanncgnaag nngangaang gngacnccg cccggcngg aaggaanngn nnnnnnnngn 1740
nnnnngnnnn nnnnnnnnnn aagcccnngn aacggcggn gaacaaacnn ccaaggagcg 1800
aaaccgcggg aagccgaccn gcacgaang ngnaangann nnnnnncgcc nnnnnnnnnc 1860
ncngngaann nannnnngna agagcnnnnn cncgcnnnnn gacggaaaga cccnngnan 1920
cacnnannnn nnangnnnnn nnnnnnnnnn gnnagnaagg nggagncnnn gannnnnnnn 1980
cgnnagnnnn nnnnggagncn nnnngnnaac nacnccnnnn nnnnnnnnnc aacnnnnnnn 2040
nnnnnancnn nnnnnngaca ngnnngnnng gnagnacgg gcggnncccc naaanngaac 2100
ggaggngnnc naaggnnncn annnnggng gnacnnnnn nnagnnaann ngnanaagnn 2160
ngcnnacggn agnnnnnacnn nncgagcagn nncgaaagn ggnnnaggac cggngggnnn 2220
nnggaagngc cncgccaacg gaaaaagnac ncnggggaaa caggcnannn ncccaagagn 2280
canacgacgg nnnngggcac ccgagcgggc ncncaccgg gcgnagnngg cccaagggnn 2340
ggcgcgccnn aaagnggnac gngagcggn anaacgcgga gacagnggc cacngnnng 2400
ngngnnngan nngannngnn ngncnagac gagaggaccg gnnngnacnn ancncgggn 2460
ncngnnngc canngcann gcngnagca nnnnggnnn gaaanngcga angcacaagn 2520
nngaancnnn cnnnnagann agnnncnncn nnnnnnnnn nnnnnnnnag nnnnnnnna 2580
gannannng ngaaggngg nngnaagnn nggnannnn nnagnnnacn nnacaannnn 2640
cnnnnnnnn nnnn 2654

```

<210> 3
 <211> 13
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Primer

<400> 3
 cgtggtgacc ctt

13

<210> 4
 <211> 14

<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 4
cgtcgtcacc gcta 14

<210> 5
<211> 13
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 5
cgtggtaccc ctt 13

<210> 6
<211> 90
<212> RNA
<213> Bacillus anthracis

<220>

<221> misc_feature
<222> 20
<223> n = A, U, C or G

<400> 6
gcgaagaacc uuaccaggun uugacauccu cugacaaccc uagagauagg gcuucuccuu 60
cgggagcaga gugacaggug gugcaugguu 90

<210> 7
<211> 90
<212> RNA
<213> Bacillus cereus

<400> 7
gcgaagaacc uuaccagguc uugacauccu cugaaaaccc uagagauagg gcuucuccuu 60
cgggagcaga gugacaggug gugcaugguu 90

<210> 8
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 8
gtgagatggt gggттаagtc ccgтаacgag 30

<210> 9
<211> 21

<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 9
gacgtcatcc ccaccttcct c 21

<210> 10
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 10
atgttggtt aagtcacgca acgag 25

<210> 11
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 11
ttgacgtcat cccaccttc ctc 23

<210> 12
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 12
ttaagtcacg caacgatcg aa 22

<210> 13
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 13
tgacgtcatc cccaccttc tc 22

<210> 14
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 14
gctacacacg tgctacaatg

20

<210> 15
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 15
cgagttgcag actgcgatcc g

21

<210> 16
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 16
aagtcggaat cgctagtaat cg

22

<210> 17
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 17
gacgggcggt gtgtacaag

19

<210> 18
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 18
tgaacgctgg tggcatgctt aacac

25

<210> 19
<211> 22
<212> DNA
<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 19

tacgcattac tcacccgtcc gc

22

<210> 20

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 20

gtggcatgcc taatacatgc aagtcg

26

<210> 21

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 21

ttactcaccg gtcgcgcgt

20

<210> 22

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 22

taacacatgc aagtcgaacg

20

<210> 23

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 23

ttactcaccg gtcgcgc

17

<210> 24

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 24
gtgtagcggt gaaatgcg 18

<210> 25
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 25
gtatctaatac ctgtttgctc cc 22

<210> 26
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 26
agaacaccga tggcgaaggc 20

<210> 27
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 27
cgtggactac cagggtatct a 21

<210> 28
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 28
ggattagaga ccctggtagt cc 22

<210> 29
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 29
ggccgtactc cccaggcg 18

<210> 30
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 30
ggattagata ccctggtagt ccacgc

26

<210> 31
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 31
ggccgtactc cccaggcg

18

<210> 32
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 32
tagataccct ggtagtccac gc

22

<210> 33
<211> 15
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 33
cgtactcccc aggcg

15

<210> 34
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 34
ttcgaatgcaa cgcgaagaac ct

22

<210> 35

<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 35
acgagctgac gacagccatg

20

<210> 36
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 36
acgcgaagaa ccttacc

17

<210> 37
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 37
acgacacgag ctgacgac

18

<210> 38
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 38
ctgacacctg cccggtgc

18

<210> 39
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 39
gaccgttata gttacggcc

19

<210> 40
<211> 25
<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 40

tctgtcccta gtacgagagg accgg

25

<210> 41

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 41

tgcttagatg ctttcagc

18

<210> 42

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 42

ctgtccctag tacgagagga ccgg

24

<210> 43

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 43

gtttcatgct tagatgcttt cagc

24

<210> 44

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 44

ggggagtgaag agagatcctg aaacgg

26

<210> 45

<211> 21

<212> DNA

<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 45
acaaaaggta cgccgtcacc c 21

<210> 46
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 46
ggggagtgaag agagatcctg aaaccg 26

<210> 47
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 47
acaaaaggca cgccatcacc c 21

<210> 48
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 48
cgagaggggaa acaaccacga cc 22

<210> 49
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 49
tggtgtcttc taagccaac 19

<210> 50
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 50
tgctcgtggt gcacaagtaa cggatatta

29

<210> 51
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 51
tgctgctttc gcatggtaa ttgcttcaa

29

<210> 52
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 52
caaaacttat taggtaagcg tggtgact

28

<210> 53
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 53
tcaagcgcca tttcttttgg taaaccacat

30

<210> 54
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 54
caaaacttat taggtaagcg tggtgact

28

<210> 55
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 55

tcaagcgcca tctctttcgg taatccacat

30

<210> 56

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 56

taagaagccg gaaaccatca actaccg

27

<210> 57

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 57

ggcgcttgta cttaccgcac

20

<210> 58

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 58

tgattctggt gcccggtggt

19

<210> 59

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 59

ttggccatca ggccacgcat ac

22

<210> 60

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 60

tgattccggt gcccggtggt

19

<210> 61
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 61
ttggccatca gaccacgcat ac 22

<210> 62
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 62
ctggcaggta tgcgtggtct gatg 24

<210> 63
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 63
cgcaccgtgg gttgagatga agtac 25

<210> 64
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 64
cttgctggta tgcgtggtct gatg 24

<210> 65
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 65
cgcacatgc gtagagatga agtac 25

<210> 66
<211> 26

<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 66
cgtcgggtga ttaaccgtaa caaccg 26

<210> 67
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 67
gtttttcggt gcgtacgatg atgtc 25

<210> 68
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 68
cgtcgtgtaa ttaaccgtaa caaccg 26

<210> 69
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 69
acgtttttcg ttttgaacga taatgct 27

<210> 70
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 70
caaaggtaag caaggtcgtt tccgtca 27

<210> 71
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 71
cgaacggcct gagtagtcaa cacg 24

<210> 72
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 72
caaaggttaag caaggacgtt tccgtca 27

<210> 73
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 73
cgaacggcca gagtagtcaa cacg 24

<210> 74
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 74
tagactgccc aggacacgct g 21

<210> 75
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 75
gccgtccatc tgagcagcac c 21

<210> 76
<211> 21
<212> DNA
<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 76

ttgactgccc aggtcacgct g

21

<210> 77

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 77

gccgtccatt tgagcagcac c

21

<210> 78

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 78

aactaccgtc cgcagttcta cttcc

25

<210> 79

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 79

gttgtcgcca ggcataacca tttc

24

<210> 80

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 80

aactaccgtc ctcagtteta cttcc

25

<210> 81

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 81
gttggtcacca ggcattacca ttct 24

<210> 82
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 82
ccacagttct acttccgtac tactgacg 28

<210> 83
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 83
tccaggcatt accatttcta ctccttctgg 30

<210> 84
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 84
gacctacagt aagaggttct gtaatgaacc 30

<210> 85
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 85
tccaagtgt ggtttacccc atgg 24

<210> 86
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 86
catccacacg gtggtggtga agg 23

<210> 87
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 87
gtgctggttt accccatgga gt 22

<210> 88
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 88
cgtgttgact attcggggcg ttcag 25

<210> 89
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 89
attcaagagc catttctttt ggtaaaccac 30

<210> 90
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 90
tcaacaacct cttggaggta aagctcagt 29

<210> 91
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 91
tttcttgaag agtatgagct gctccgtaag 30

<210> 92

<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 92
catccacacg gtggtggtga agg 23

<210> 93
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 93
tgttttgtat ccaagtgtg gttacccc 29

<210> 94
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 94
cgtggcggcg tggttatcga 20

<210> 95
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 95
cggtagaac tggatgtcgc cggt 24

<210> 96
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 96
tatcgctcag gcgaactcca ac 22

<210> 97
<211> 21
<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 97

gctggattcg cctttgctac g

21

<210> 98

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 98

tgtaatgaac cctaagacc atccacacgg

30

<210> 99

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 99

ccaagtgctg gtttacccca tggagta

27

<210> 100

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 100

taatgaaccc taatgaccat ccacacggtg

30

<210> 101

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 101

tccaagtgtt gggttaccac atggag

26

<210> 102

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 102

cttgaggta agtctcattt tggtagggca

29

<210> 103

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 103

cgtataagct gcaccataag cttgtaatgc

30

<210> 104

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 104

cgacgcgtg cgcttcac

18

<210> 105

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 105

gcgttcaca gcttggtgca gaag

24

<210> 106

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 106

gaccacctcg gcaaccgt

18

<210> 107

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 107
ttcgctctcg gcctggcc 18

<210> 108
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 108
gcactatgca cacgtagatt gtcttg 27

<210> 109
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 109
tatagcacca tccatctgag cggcac 26

<210> 110
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 110
cgcggtactt caacgacagc ca 22

<210> 111
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 111
cgcggtcggc tcgttgatga 20

<210> 112
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 112

cttctgcaac aagctgtgga acgc

24

<210> 113

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 113

tcgcagttca tcagcacgaa gcg

23

<210> 114

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 114

aagacgacct gcacgggc

18

<210> 115

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 115

gcgctccacg tcttcacgc

19

<210> 116

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 116

ctgttttag tacgagagga cc

22

<210> 117

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 117

ttcgtgotta gatgetttca g

21

<210> 118
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 118
acgcgaagaa ccttacc

17

<210> 119
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 119
acgacacgag ctgacgac

18

<210> 120
<211> 14
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 120
cgaagaacct tacc

14

<210> 121
<211> 12
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 121
acacgagctg ac

12

<210> 122
<211> 14
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 122
cgaagaacct tacc

14

<210> 123
<211> 12

<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 123
acacgagctg ac 12

<210> 124
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 124
cctgataagg gtgaggtcg 19

<210> 125
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 125
acgtccttca tcgcctctga 20

<210> 126
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 126
gttgtgaggt taagcgacta ag 22

<210> 127
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 127
ctatcgggtca gtcaggagta t 21

<210> 128
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 128
g ttgtgaggt taagcgacta ag 22

<210> 129
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 129
t tgcacgcggg ttggttaagtc 20

<210> 130
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 130
a tactcctga ctgaccgata g 21

<210> 131
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 131
a acatagcct tctccgtcc 19

<210> 132
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 132
g acttaccaa cccgatgcaa 20

<210> 133
<211> 24
<212> DNA
<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 133

taccttagga ccgttatagt tacg

24

<210> 134

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 134

ggacggagaa ggctatggt

19

<210> 135

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 135

ccaaacaccg ccgtcgatat

20

<210> 136

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 136

cgtaactata acggtcctaa ggta

24

<210> 137

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 137

gcttacacac ccggcctatc

20

<210> 138

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 138
atatcgacgg cggtgtttgg 20

<210> 139
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 139
gcgtgacagg caggtattc 19

<210> 140
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 140
agtctcaaga gtgaacacgt aa 22

<210> 141
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 141
gctgctggca cggagtta 18

<210> 142
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 142
gacacggtcc agactcctac 20

<210> 143
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 143
ccatgcagca cctgtctc 18

<210> 144
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 144
gatctggagg aataccggtg

20

<210> 145
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 145
acggttacct tggtacgact

20

<210> 146
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 146
gagagcaagc ggacctcata

20

<210> 147
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 147
cctcctgcgt gcaaagc

17

<210> 148
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 148
tggaagatct gggtcaggc

19

<210> 149

<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 149
caatctgctg acggatctga gc 22

<210> 150
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 150
gtcgtgaaaa cgagctggaa ga 22

<210> 151
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 151
catgatggtc acaaccgg 18

<210> 152
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 152
tggcgaacct ggtgaacgaa gc 22

<210> 153
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 153
ctttcgcttt ctcgaactca accat 25

<210> 154
<211> 26
<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 154

cgtcagggtg aattccgtga agttaa

26

<210> 155

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 155

aacttcgcct tcgggtcatgt t

21

<210> 156

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 156

ggtgaaagaa gttgcctcta aagc

24

<210> 157

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 157

ttcaggtcca tcgggttcat gcc

23

<210> 158

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 158

cgtggcggcg tggttatcga

20

<210> 159

<211> 20

<212> DNA

<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 159
acgaactgga tgcgccgtt 20

<210> 160
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 160
cggaattact gggcgtaaag 20

<210> 161
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 161
cgcatctcac cgctacac 18

<210> 162
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 162
accagtgct gctgaaccgt gc 22

<210> 163
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 163
gttcaaagtc ctggataccc a 21

<210> 164
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 164
gggagcaaac aggattagat ac 22

<210> 165
<211> 15
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 165
cgtactcccc aggcg 15

<210> 166
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 166
tggcccgaag gaagctgagc g 21

<210> 167
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 167
acgcgggcat gcagagatgc c 21

<210> 168
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 168
atgttggtt aagtcgagc 19

<210> 169
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 169

tgacgtcatc cccaccttcc

20

<210> 170

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 170

cttgtacaca ccgcccgtc

19

<210> 171

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 171

aaggaggtga tccagcc

17

<210> 172

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 172

cggattggag tctgcaactc g

21

<210> 173

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 173

gacgggcggt gtgtacaag

19

<210> 174

<211> 15

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 174

ggtggatgcc ttggc

15

<210> 175
<211> 16
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 175
gggtttcccc attcgg

16

<210> 176
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 176
gggaactgaa acatctaagt a

21

<210> 177
<211> 15
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 177
ttcgtctgcc gctac

15

<210> 178
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 178
taccctaaac cgacacagg

19

<210> 179
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 179
ccttctcccg aagttacg

18

<210> 180
<211> 19

<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 180
ccgtaacttc gggagaagg 19

<210> 181
<211> 15
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 181
caccgggcag gcgtc 15

<210> 182
<211> 16
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 182
gacgcctgcc cgggtgc 16

<210> 183
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 183
ccgacaagga atttcgctac c 21

<210> 184
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 184
aagggtactcc ggggataaca ggc 23

<210> 185
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 185
agccgacatc gaggtgccaa ac 22

<210> 186
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 186
gacagtccgg tccctatc 18

<210> 187
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 187
ccggtcctct cgtacta 17

<210> 188
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 188
tagtacgaga ggaccgg 17

<210> 189
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 189
ttagatgctt tcagcactta tc 22

<210> 190
<211> 24
<212> DNA
<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 190

aaactagata acagtagaca tcac

24

<210> 191

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 191

gtgcgccctt tctaactt

18

<210> 192

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 192

agagtttgat catggctcag

20

<210> 193

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 193

actgctgcct cccgtag

17

<210> 194

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 194

cactggaact gagacacgg

19

<210> 195

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 195
ctttacgccc agtaattccg 20

<210> 196
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 196
ccagcagccg cggtaatac 19

<210> 197
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 197
gtatctaatac ctgtttgctc cc 22

<210> 198
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 198
gtgtagcggg gaaatgcg 18

<210> 199
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 199
ggtaagggtt ttgcggttg 19

<210> 200
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 200
aagcgggtgga gcatgtgg 18

<210> 201
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 201
attgtagcac gtgtgtagcc c 21

<210> 202
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 202
caagtcacatca tggccctta 19

<210> 203
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 203
aaggaggtga tccagcc 17

<210> 204
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 204
agagtttgat catggctcag 20

<210> 205
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 205
aaggaggtga tccagcc 17

<210> 206

<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 206
acctgcccag tgctggaag 19

<210> 207
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 207
tcgtacett aggaccgt 18

<210> 208
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 208
gccttgta caacctccgt c 21

<210> 209
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 209
cacggctacc ttgttacgac 20

<210> 210
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 210
ttgtacacac cgcctgcac ac 22

<210> 211
<211> 20
<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 211

ccttggtacg acttcacccc

20

<210> 212

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 212

tacggtgaat acgttcccg g

21

<210> 213

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 213

accttggtac gaattcaccc ca

22

<210> 214

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 214

accacgccgt aaacgatga

19

<210> 215

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 215

cccccgtaa ttcctttgag t

21

<210> 216

<211> 22

<212> DNA

<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 216
gataccctgg tagtccacac cg 22

<210> 217
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 217
gccttgcgac cgtactccc 19

<210> 218
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 218
tagataccct ggtagtccac gc 22

<210> 219
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 219
gcgaccgtac tccccagg 18

<210> 220
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 220
tagtcccgcga acgagcgc 18

<210> 221
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 221
gacgtcatcc ccaccttcct cc 22

<210> 222
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 222
tagaacgtcg cgagacagtt cg 22

<210> 223
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 223
agtccatccc ggtcctctcg 20

<210> 224
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 224
gaggaaagtc cgggctc 17

<210> 225
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 225
ataagccggg ttctgtcg 18

<210> 226
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 226

gaggaaagtc catgctcgc 19

<210> 227
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 227
gtaagccatg ttttgttcca tc 22

<210> 228
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 228
gaggaaagtc cgggctc 17

<210> 229
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 229
ataagccggg ttctgtcg 18

<210> 230
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 230
gcgggatacct ctagaggtgt taaatagcct ggcag 35

<210> 231
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 231
gcgggatacct ctagaagacc tcctgcgtgc aaagc 35

<210> 232
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 232
gaggaaagtc catgctcac 19

<210> 233
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 233
ataagccatg ttctgttcca tc 22

<210> 234
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 234
atgttgggtt aagtccgc 19

<210> 235
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 235
aaggaggtga tccagcc 17

<210> 236
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 236
cggaattact gggcgtaaag 20

<210> 237
<211> 22

<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 237
gtatctaatac ctgtttgctc cc

22

<210> 238
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 238
atgttgggtt aagtccgc

19

<210> 239
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 239
tgacgtcatg cccaccttcc

20

<210> 240
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 240
atgttgggtt aagtccgc

19

<210> 241
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 241
tgacgtcatg gccaccttcc

20

<210> 242
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 242
gcgggatcct ctagacctga taagggtgag gtcg 34

<210> 243
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 243
gcgggatcct ctagagcgtg acaggcaggt attc 34

<210> 244
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 244
acgcgaagaa ccttacc 17

<210> 245
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 245
gacgggcggt gtgtacaag 19

<210> 246
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer.

<400> 246
gtgtagcggg gaaatgcg 18

<210> 247
<211> 21
<212> DNA
<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 247
cgagttgcag actgcgatcc g

21

<210> 248
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 248
taacacatgc aagtcgaacg

20

<210> 249
<211> 15
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 249
cgtactcccc aggcg

15

<210> 250
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 250
taacacatgc aagtcgaacg

20

<210> 251
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 251
acgacacgag ctgacgac

18

<210> 252
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 252
acaacgaagt acaatacaag ac 22

<210> 253
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 253
cttctacatt tttagccatc ac 22

<210> 254
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 254
ttaagtcccg caacgagcgc aa 22

<210> 255
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 255
tgacgtcatc cccaccttcc tc 22

<210> 256
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 256
tgagtgatga aggccttagg gttgtaaa 28

<210> 257
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 257
cggctgctgg cacgaagtta g 21

<210> 258
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 258
atggacaagg ttggcaagga agg

23

<210> 259
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 259
tagccgcggt cgaattgcat

20

<210> 260
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 260
aaggaaggcg tgatcaccgt tgaaga

26

<210> 261
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 261
ccgcggtcga attgcatgcc ttc

23

<210> 262
<211> 16
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 262
acgcgctgcg cttcac

16

<210> 263

<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 263
ttgcagaagt tgcggtagcc

20

<210> 264
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 264
tcgaccacct gggcaacc

18

<210> 265
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 265
atcaggtcgt gcggcatca

19

<210> 266
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 266
cacggtgccg gcgtact

17

<210> 267
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 267
gcggtcggct cgttgatgat

20

<210> 268
<211> 25
<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 268

ttggaggtaa gtctcat ttt ggtgg

25

<210> 269

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 269

aagctgcacc ataagcttgt aatgc

25

<210> 270

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 270

cagcgtttcg gcgaaatgga

20

<210> 271

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 271

cgacttgacg gttaacattt cctg

24

<210> 272

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 272

gggcagcgtt tcggcgaaat gga

23

<210> 273

<211> 27

<212> DNA

<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 273
gtccgacttg acggtcaaca ttctctg 27

<210> 274
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 274
caggagtcgt tcaactcgat ctacatgat 29

<210> 275
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 275
acgccatcag gccacgcat 19

<210> 276
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 276
gcacaacctg cggtgcg 18

<210> 277
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 277
acggcacgag gtagtcgc 18

<210> 278
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

tggtactcac ccgtctgccca ct 22

<210> 284

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 284

accgagcaag gagaccagc 19

<210> 285

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 285

tataacgcac atcgtcaggg tga 23

<210> 286

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 286

agaccaatt acattggctt 20

<210> 287

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 287

ccagtgtgt tgtagtacat 20

<210> 288

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 288

atgtactaca acagtactgg 20

<210> 289
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 289
caagtcaacc acagcattca 20

<210> 290
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 290
gggcttatgt actacaacag 20

<210> 291
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 291
tctgtcttgc aagtcaacca c 21

<210> 292
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 292
ggaatttttt gatggtagag a 21

<210> 293
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 293
taaagcaciaa tttcaggcg 19

<210> 294
<211> 20

<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 294
tagatctggc tttctttgac 20

<210> 295
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 295
atatgagtat ctggagtctg c 21

<210> 296
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 296
ggaaagacat tactgcagac a 21

<210> 297
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 297
ccaacttgag gctctggctg 20

<210> 298
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 298
acagacactt accagggctg 19

<210> 299
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 299

actgtggtgt catctttgtc

. 20

<210> 300

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 300

tcactaaaga caaaggtctt cc

22

<210> 301

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 301

ggcttcgccg tctgtaattt c

21

<210> 302

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 302

oggatccaag ctaatctttg g

21

<210> 303

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 303

ggtatgtact cataggtggtt ggtg

24

<210> 304

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 304

agaccaatt acattggctt

20

<210> 305

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 305

ccagtgtgt tgtagtacat

20

<210> 306

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 306

atgtactaca acagtactgg

20

<210> 307

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 307

caagtcaacc acagcattca

20

<210> 308

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 308

gggcttatgt actacaacag

20

<210> 309

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 309
tctgtcttgc aagtcaacca c 21

<210> 310
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 310
ggaatttttt gatggtagag a 21

<210> 311
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 311
taaagcacaa tttcaggcg 19

<210> 312
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 312
tagatctggc tttctttgac 20

<210> 313
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 313
atatgagtat ctggagtctg c 21

<210> 314
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 314
cggatccaag ctaatctttg g 21

<210> 315
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 315
ggtatgtact cataggtggt ggtg

24

<210> 316
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 316
aacagaccca attacattgg ctt

23

<210> 317
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 317
gaggcacttg tatgtggaaa gg

22

<210> 318
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 318
atgcctaaca gaccgaatta cat

23

<210> 319
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 319
ttcatgtagt cgtaggtggt gg

22

<210> 320

<220>
<223> PCR Primer

<400> 330
ggtcgttatg tgcctttcca cat

<210> 331
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 331
tcctttctga agttccactc atagg

<210> 332
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 332
acaacattgg ctaccagggc tt

<210> 333
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 333
cctgcctgct cataggctgg aagtt

<210> 334
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 334
ggattagaga ccctggtagt cc

<210> 335
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

23

25

22

25

22

<400> 335
ggccgtactc cccaggcg 18

<210> 336
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 336
ttcgatgcaa cgcgaagaac ct 22

<210> 337
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 337
acgagctgac gacagccatg 20

<210> 338
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 338
tctgtcccta gtacgagagg accgg 25

<210> 339
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 339
tgcttagatg ctttcagc 18

<210> 340
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 340

ctggcaggta tgcgtggtct gatg 24

<210> 341
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 341
cgcacccgtgg gttgagatga agtac 25

<210> 342
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 342
ggggattcag ccatcaaagc agctattgac 30

<210> 343
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 343
ccaacctttt ccacaacaga atcagc 26

<210> 344
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 344
ccttacttcg aactatgaat cttttggaag 30

<210> 345
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 345
cccatttttt cacgcatgct gaaaatatc 29

<210> 346
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 346
cgcaaaaaaa tccagctatt agc 23

<210> 347
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 347
aaactatttt tttagctata ctcgaacac 29

<210> 348
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 348
atgattacaa ttcaagaagg tcgtcacgc 29

<210> 349
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 349
ttggacctgt aatcagctga atactgg 27

<210> 350
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 350
gatgactttt tagctaattgg tcaggcagc 29

<210> 351
<211> 29

<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 351
aatcgacgac catcttgga agatttctc 29

<210> 352
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 352
gcttcaggaa tcaatgatgg agcag 25

<210> 353
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 353
gggtctacac ctgcacttgc ataac 25

<210> 354
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 354
gtactgaatc cgcctaag 18

<210> 355
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 355
gtgaataaag tatcgcccta ata . 23

<210> 356
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 356
gaagttgaac cgggatca 18

<210> 357
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 357
attatcggtc gttgttaatg t 21

<210> 358
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 358
ctgtctgtag ataaactagg att 23

<210> 359
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 359
cggtcttctc tggaggat 18

<210> 360
<211> 15
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 360
cgatactacg gacgc 15

<210> 361
<211> 22
<212> DNA
<213> Artificial Sequence

<220>

<210> 372
<211> 15
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 372
cgatactacg gacgc

15

<210> 373
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 373
ctttatgaat tactttacat at

22

<210> 374
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 374
ctcctccatc actaggaa

18

<210> 375
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 375
ctataacatt caaagcttat tg

22

<210> 376
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 376
cgcgataata gatagtgcata aac

23

<210> 377

<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primers

<400> 377
gcttcacca ggtcattaa